

Mark Scheme (Results)

June 2011

GCE Core Mathematics C3 (6665) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025 or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link: http://www.edexcel.com/Aboutus/contact-us/

June 2011
Publications Code UA027660
All the material in this publication is copyright
© Edexcel Ltd 2011



EDEXCEL GCE MATHEMATICS

General Instructions for Marking

- 1. The total number of marks for the paper is 75.
- 2. The Edexcel Mathematics mark schemes use the following types of marks:
 - M marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - B marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod benefit of doubt
- ft follow through
- the symbol wi√l be used for correct ft
- cao correct answer only
- cso correct solution only. There must be no errors in this part of the question to obtain this mark
- isw ignore subsequent working
- awrt answers which round to
- SC: special case
- oe or equivalent (and appropriate)
- dep dependent
- indep independent
- · dp decimal places
- sf significant figures
- * The answer is printed on the paper
- Lather the second mark is dependent on gaining the first mark

Question Number	Scheme	Marks	
1 (a)	$\frac{1}{(x^2+3x+5)} \times \dots , = \frac{2x+3}{(x^2+3x+5)}$	M1,A1 (2	5)
(b)	Applying $\frac{vu'-uv'}{v^2}$ $\frac{x^2 \times -\sin x - \cos x \times 2x}{(x^2)^2} = \frac{-x^2 \sin x - 2x \cos x}{x^4} = \frac{-x \sin x - 2\cos x}{x^3} \text{ oe}$	M1, A2,1,0 (3 5 Mark	
2 (a)	f(0.75) = -0.18 $f(0.85) = 0.17$ Change of sign, hence root between x=0.75 and x=0.85	M1 A1 (2	2)
(b)	Sub $x_0=0.8$ into $x_{n+1} = [\arcsin(1-0.5x_n)]^{\frac{1}{2}}$ to obtain x_1 Awrt $x_1=0.80219$ and $x_2=0.80133$ Awrt $x_3=0.80167$	M1 A1 A1	
(c)	$f(0.801565) = -2.7 \times 10^{-5}$ $f(0.801575) = +8.6 \times 10^{-6}$	(3 M1A1	
	Change of sign and conclusion See Notes for continued iteration method	A1 (3	5)
		8 Mark	.S

Question	Scheme	Marks
Number 3 (a)	A.Y	
S (a)	V shape vertex on y axis &both branches of graph cross x axis 'y' co-ordinate of R is -6	B1 B1 B1
	(0,-6)	(3)
(b)	(-4,3) W shape 2 vertices on the negative x axis. W in both quad 1 & quad 2. $R'=(-4,3)$	B1 B1dep B1 (3) 6 Marks
4 (a)	$y = 4 - \ln(x + 2)$ $\ln(x + 2) = 4 - y$ $x + 2 = e^{4-y}$ $x = e^{4-y} - 2$ $f^{-1}(x) = e^{4-x} - 2$ oe	M1 M1A1 (3)
(b)	$x \le 4$	B1 (1)
(c)	$fg(x) = 4 - \ln(e^{x^2} - 2 + 2)$	M1
	$fg(x) = 4 - x^2$	dM1A1 (3)
(d)	$fg(x) \le 4$	B1ft (1)
		8 Marks

Question Number	Scheme	Marks
5 (a)	p=7.5	B1 (1)
(b)	$2.5 = 7.5e^{-4k}$	(1) M1
	$e^{-4k} = \frac{1}{3}$	M1
	$-4k = \ln(\frac{1}{3})$	dM1
	$-4k = -\ln(3)$ $k = \frac{1}{4}\ln(3)$	
	$k = \frac{1}{4} \ln(3)$	A1*
	See notes for additional correct solutions and the last A1	(4)
(c)	dm	
(0)	$\frac{dm}{dt} = -kpe^{-kt}$ ft on their p and k	M1A1ft
	$-\frac{1}{4}\ln 3 \times 7.5e^{-\frac{1}{4}(\ln 3)t} = -0.6\ln 3$	
	$e^{-\frac{1}{4}(ln3)t} = \frac{2.4}{7.5} = (0.32)$	M1A1
	$-\frac{1}{4}(\ln 3)t = \ln(0.32)$	dM1
	<i>t</i> =4.1486 4.15 or awrt 4.1	A1
		(6)
		11Marks

Question Number	Scheme		Marks
6 (a)	$\frac{1}{\sin 2\theta} - \frac{\cos 2\theta}{\sin 2\theta} = \frac{1 - \cos 2\theta}{\sin 2\theta}$		M1
	$=\frac{2\sin^2\theta}{2\sin\theta\cos\theta}$		M1A1
	$=\frac{\sin\theta}{\cos\theta}=\tan\theta$	cso	A1* (4)
(b)(i)	$\tan 15^\circ = \frac{1}{\sin 30^\circ} - \frac{\cos 30^\circ}{\sin 30^\circ}$		M1
	$\tan 15^{\circ} = \frac{\frac{1}{1}}{\frac{1}{2}} - \frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}} = 2 - \sqrt{3}$	cso	dM1 A1*
(b)(ii)	$\tan 2x = 1$		M1
	$2x = 45^{\circ}$		A1
	$2x = 45^{\circ} + 180^{\circ}$		M1
	$x = 22.5^{\circ}, 112.5^{\circ}, 202.5^{\circ}, 292.5^{\circ}$		A1(any two) A1 (5)
	Alt for (b)(i) $\tan 15^{\circ} = \tan(60^{\circ} - 45^{\circ})$ or $\tan(45^{\circ} - 30^{\circ})$		12 Marks
	$\tan 15^{\circ} = \frac{\tan 60 - \tan 45}{1 + \tan 60 \tan 45} \text{ or } \frac{\tan 45 - \tan 30}{1 + \tan 45 \tan 30}$		M1
	$\tan 15^{\circ} = \frac{\sqrt{3} - 1}{1 + \sqrt{3}} \text{ or } \frac{1 - \frac{\sqrt{3}}{3}}{1 + \frac{\sqrt{3}}{3}}$		M1
	Rationalises to produce $tan15^{\circ} = 2 - \sqrt{3}$		A1*

Question Number	Scheme	Marks
7 (a)	$x^{2} - 9 = (x+3)(x-3)$ $\frac{4x-5}{(2x+1)(x-3)} - \frac{2x}{(x+3)(x-3)}$	B1
	$= \frac{(4x-5)(x+3)}{(2x+1)(x-3)(x+3)} - \frac{2x(2x+1)}{(2x+1)(x+3)(x-3)}$	M1
	$=\frac{5x-15}{(2x+1)(x-3)(x+3)}$	M1A1
	$= \frac{5(x-3)}{(2x+1)(x-3)(x+3)} = \frac{5}{(2x+1)(x+3)}$	A1*
		(3)
(b)	$f(x) = \frac{5}{2x^2 + 7x + 3}$	
	$f'(x) = \frac{-5(4x+7)}{(2x^2+7x+3)^2}$	M1 M1 A1
	$f'\left(-1\right) = -\frac{15}{4}$	M1A1
	Uses m_1m_2 =-1 to give gradient of normal= $\frac{4}{15}$	M1
	$\frac{y - (-\frac{5}{2})}{(x1)} = their \frac{4}{15}$	M1
	$y + \frac{5}{2} = \frac{4}{15}(x+1)$ or any equivalent form	A1
		(8)
		13 Marks

Question	Scheme	Marks
Number 8		
(a)	$R^2 = 2^2 + 3^2$ $R = \sqrt{13} \text{ or } 3.61 \dots$	M1 A1
	$\tan \alpha = \frac{3}{2}$	M1
	$\alpha = 0.983$	A1
		(4)
(b)	$f'(x) = 2e^{2x}\cos 3x - 3e^{2x}\sin 3x$	M1A1A1
	$=e^{2x}(2\cos 3x - 3\sin 3x)$	M1
	$=e^{2x}(R\cos(3x+\alpha)$	
	$= Re^{2x}\cos(3x + \alpha)$	A1* cso
		(5)
		M1
(c)	$f'(x) = 0 \Rightarrow \cos(3x + \alpha) = 0$	
	$3x + \alpha = \frac{\pi}{2}$	M1
	0.106	0.20
	<i>x</i> =0.196 awrt	
		(3)
		12 Marks
	Alternative to part (c) \Rightarrow	
	$f'(x) = 0 \Rightarrow 2\cos 3x - 3\sin 3x = 0$	M1
	,	
	$\tan 3x = \frac{2}{3}$	M1
	x=0.196 awr	t 0.20 A1
		(3)
		(3)

Further copies of this publication are available from Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email <u>publication.orders@edexcel.com</u>

Order Code UA027660 June 2011

For more information on Edexcel qualifications, please visit www.edexcel.com/quals

Pearson Education Limited. Registered company number 872828 with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE





