



GCSE

Mathematics

Unit **J560/06**: Higher Tier Paper 6

General Certificate of Secondary Education

Mark Scheme for November 2017

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2017

1. Annotations used in the detailed Mark Scheme.

Annotation	Meaning
✓	Correct
✗	Incorrect
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working (after correct answer obtained), provided method has been completed
M0	Method mark awarded 0
M1	Method mark awarded 1
M2	Method mark awarded 2
A1	Accuracy mark awarded 1
B1	Independent mark awarded 1
B2	Independent mark awarded 2
MR	Misread
SC	Special case
^	Omission sign

These should be used whenever appropriate during your marking.

The **M**, **A**, **B** etc annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate these scripts to show how the marks have been awarded. It is not mandatory to use annotations for any other marking, though you may wish to use them in some circumstances.

Subject-Specific Marking Instructions

2. **M** marks are for using a correct method and are not lost for purely numerical errors.
A marks are for an accurate answer and depend on preceding **M** (method) marks. Therefore **M0 A1** cannot be awarded.
B marks are independent of **M** (method) marks and are for a correct final answer, a partially correct answer, or a correct intermediate stage.
SC marks are for special cases that are worthy of some credit.

3. Unless the answer and marks columns of the mark scheme specify **M** and **A** marks etc, or the mark scheme is 'banded', then if the correct answer is clearly given and is not from wrong working **full marks** should be awarded.

Do not award the marks if the answer was obtained from an incorrect method, ie incorrect working is seen and the correct answer clearly follows from it.

4. Where follow through (**FT**) is indicated in the mark scheme, marks can be awarded where the candidate's work follows correctly from a previous answer whether or not it was correct.

Figures or expressions that are being followed through are sometimes encompassed by single quotation marks after the word *their* for clarity, eg FT $180 \times (\textit{their} '37' + 16)$, or FT $300 - \sqrt{(\textit{their} '5^2 + 7^2')}$. Answers to part questions which are being followed through are indicated by eg FT $3 \times \textit{their} (a)$.

For questions with FT available you must ensure that you refer back to the relevant previous answer. You may find it easier to mark these questions candidate by candidate rather than question by question.

5. Where dependent (**dep**) marks are indicated in the mark scheme, you must check that the candidate has met all the criteria specified for the mark to be awarded.
6. The following abbreviations are commonly found in GCSE Mathematics mark schemes.

- **cao** means **correct answer only**.
- **figs 237**, for example, means any answer with only these digits. You should ignore leading or trailing zeros and any decimal point eg 237000, 2.37, 2.370, 0.00237 would be acceptable but 23070 or 2374 would not.
- **isw** means **ignore subsequent working** (after correct answer obtained).
- **nfww** means **not from wrong working**.
- **oe** means **or equivalent**.
- **rot** means **rounded or truncated**.
- **seen** means that you should award the mark if that number/expression is seen anywhere in the answer space, including the answer line,
even if it is not in the method leading to the final answer.
- **soi** means **seen or implied**.

7. Make no deductions for wrong work after an acceptable answer unless the mark scheme says otherwise, indicated for example by the instruction 'mark final answer'.

8. As a general principle, if two or more methods are offered, mark only the method that leads to the answer on the answer line. If two (or more) answers are offered, mark the poorer (poorest).
9. When the data of a question is consistently misread in such a way as not to alter the nature or difficulty of the question, please follow the candidate's work and allow follow through for **A** and **B** marks. Deduct 1 mark from any **A** or **B** marks earned and record this by using the MR annotation. **M** marks are not deducted for misreads.
10. Unless the question asks for an answer to a specific degree of accuracy, always mark at the greatest number of significant figures even if this is rounded or truncated on the answer line. For example, an answer in the mark scheme is 15.75, which is seen in the working. The candidate then rounds or truncates this to 15.8, 15 or 16 on the answer line. Allow full marks for the 15.75.
11. If the correct answer is seen in the body and the answer given in the answer space is a clear transcription error allow full marks unless the mark scheme says 'mark final answer' or 'cao'. Place the annotation ✓ next to the correct answer.

If the answer space is blank but the correct answer is seen in the body allow full marks. Place the annotation ✓ next to the correct answer.

If the correct answer is seen in the working but a completely different answer is seen in the answer space, then accuracy marks for the answer are lost. Method marks would still be awarded. Use the M0, M1, M2 annotations as appropriate and place the annotation ✘ next to the wrong answer.

12. Ranges of answers given in the mark scheme are always inclusive.
13. For methods not provided for in the mark scheme give as far as possible equivalent marks for equivalent work. If in doubt, consult your Team Leader.
14. Anything in the mark scheme which is in square brackets [...] is not required for the mark to be earned, but if present it must be correct.

Question		Answer	Marks	Part marks and guidance	
1	(a)	200	2	B1 for 50 or 150 soi	Eg. answer 500 or 275 with (5×10) + ... seen
	(b)	$a = \frac{2(s-ut)}{t^2}$ oe	2	M1 for $s - ut = \frac{1}{2} at^2$	
2		30 November	3	B2 for 21 identified as LCM or answer 30 OR M1 for listing at least four multiples of 3 and at least three multiples of 7 OR M1 for listing/identifying at least four dates for running and at least three dates for cycling	3, 6, 9, 12, 15, 18, 21, ... 7, 14, 21, ... 12, 15, 18, 21, ... 16, 23, 30, ...
3	(a)	5.34	4	B1 for 1.5, 4.5, 7.5, 10.5, 13.5 M1FT for 1.5×6 4.5×10 7.5×6 10.5×2 13.5×1 soi 9, 45, 45, 21, 13.5 or 133.5 M1 for <i>their</i> $133.5 \div 25$	At least 4 midpoints correct FT midpoints or either end of range values consistently used Allow one numerical error Four correct products or 133.5 imply B1 and M1
	(b)	Exact times for each customer are not recorded oe	1		Do not accept, "Because the midpoint is used" or comments on the method used.

Question		Answer	Marks	Part marks and guidance	
4	(a)	$\frac{12}{28} = \frac{3}{7}$ or $\frac{3}{7}$ of 28 = 12 or $12 \div 28 = \frac{3}{7}$	1		Alternative $12 \div 28 = 0.428571\dots$ and $3 \div 7 = 0.428571\dots$
	(b)	Integer from 23000 to 23334	2	M1 for $10\,000 \div \frac{3}{7}$ oe If M0 then SC1 for figs 2333... seen	Accept integer from 23000 to 24000 after M1
	(c)	The growing conditions on the farm may be different to the garden oe or Sample too small oe	1		Mere reference to factors that affect growth is insufficient
5	(a)	42	2	M1 for $\frac{1.47 \times 10^7}{3.5 \times 10^5}$ oe If 0 scored SC1 for figs 42 in answer	Eg. $\frac{14\,700\,000}{350\,000}$
	(b)	$4.2[3\dots] \times 10^9$	3	B2 for 4 233 600 000 oe as answer or M1 for <i>their</i> $1.47 \times 10^7 \times 288$ If 0 scored SC1 for figs 423[...] in answer	Eg. $423.[36] \times 10^7$ <i>their</i> 1.47×10^7 converted from info in (a)

Question		Answer	Marks	Part marks and guidance										
	(c) (i)	6450	3	B2 for 6447 to 6448 or M1 for $\frac{1.47 \times 10^7}{(152 \times 15)}$ oe or figs 6447 in answer	May be in stages. NB: $152 \times 15 = 2280$									
	(ii)	Each machine makes the same amount of sweets or There are no breakdowns oe or Machines running at same rate oe or All machines run for the same time oe	1											
6	(a) (i)	$\frac{1}{5}$ of Bag A's counters [are red] or The ratio of red to yellow in Bag B is 1:3	1	Accept $1 : 4 = \frac{1}{5}$ Accept $\frac{1}{4} = 1 : 3$	Equivalentents may be percentages or decimals Eg. Bag A: 20% red, Bag B: 25% red.									
	(ii)	Correct answer is any integer multiple of this. <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th></th> <th>Red</th> <th>Yellow</th> </tr> </thead> <tbody> <tr> <td>Bag A</td> <td>4</td> <td>16</td> </tr> <tr> <td>Bag B</td> <td>5</td> <td>15</td> </tr> </tbody> </table>		Red	Yellow	Bag A	4	16	Bag B	5	15	3	B1 for (Bag A) yellow = 4 × red <i>and</i> A total = B total B1 for (Bag B) yellow = 3 × red If 0 scored SC2 for correct figures but transposed horizontally	8 32 10 30
	Red	Yellow												
Bag A	4	16												
Bag B	5	15												

Question		Answer	Marks	Part marks and guidance	
	(b)	20 nfw	3	<p>B1 for two ratios equivalent to 3:4</p> <p>M1 for <i>their</i> 15:20 reduced to (15-3):20</p> <p><u>Alternative approach</u></p> <p>B1 for two fractions equivalent to $\frac{3}{7}$</p> <p>M1 for <i>their</i> $\frac{15}{35}$ reduced to $\frac{15-3}{32}$</p>	<p>6:8, 9:12, 12:16, 15:20,...</p> <p><i>their</i> 15:20 any ratio but not 3:4</p> <p>using equivalent fractions:</p> <p>Eg $\frac{6}{14}$ or $\frac{9}{21}$ or $\frac{12}{28}$ or $\frac{15}{35}$</p> <p><i>their</i> $\frac{15}{35}$ any fraction but not $\frac{3}{7}$</p>
7		3.5	4	<p>M3 for $\frac{(629.20 - 520) \div 6}{520} [\times 100]$</p> <p>OR</p> <p>B2 for 18.2[0] seen</p> <p>OR</p> <p>B1 for 109.2[0] seen and M1 for <i>their</i> $109.2[0] \div 520 [\times 100]$</p>	

Question		Answer	Marks	Part marks and guidance
8		13.7	5	<p>M4 for $\frac{45}{360} \times 2 \times \pi \times 6 + \frac{45}{360} \times 2 \times \pi \times 2.5 + 2 \times 3.5$ oe soi by 13.67... to 13.68</p> <p>OR</p> <p>M3 for $\frac{45}{360} \times 2 \times \pi \times 6$ oe and $\frac{45}{360} \times 2 \times \pi \times 2.5$ oe soi by 4.71... and 1.96... or by 6.67... to 6.68</p> <p>OR</p> <p>M2 for $\frac{45}{360} \times 2 \times \pi \times 6$ oe or $\frac{45}{360} \times 2 \times \pi \times 2.5$ oe soi or $2 \times \pi \times 6 + 2 \times \pi \times 2.5$ oe soi by 4.71... or 1.96...or 53.4..</p> <p>OR</p> <p>M1 for $2 \times \pi \times 6$ oe or $2 \times \pi \times 2.5$ oe soi by 37.699... to 37.7 or 15.7...</p> <p>If 0 scored SC2 for $\frac{45}{360} \times \pi \times 6^2$ oe and $\frac{45}{360} \times \pi \times 2.5^2$ oe soi by 14.1... and 2.45... or by 11.6 to 11.7</p> <p>OR</p> <p>SC1 for $\frac{45}{360} \times \pi \times 6^2$ oe or $\frac{45}{360} \times \pi \times 2.5^2$ oe soi by 14.1... or 2.45...</p> <p>Method marks may be awarded for multiples of π seen in correct working. Eg. $\frac{45}{360} \times 2 \times \pi \times 6 = \frac{3}{2} \pi$ $\frac{45}{360} \times 2 \times \pi \times 2.5 = \frac{5}{8} \pi$</p>

Question		Answer	Marks	Part marks and guidance	
9		216	4	<p>M1 for $5b = 180$ oe</p> <p>and</p> <p>B1 for $[b =] 36$</p> <p>and</p> <p>M1 for <i>their</i> $36 + 180$ or <i>their</i> 36×6 provided <i>their</i> answer would be between 180 and 270</p>	<p>Eg. $b + 180 = 6b$</p> <p>soi by final answer</p>
10		<p>71 000 000 to 89 000 000 in figs or words</p> <p>people/year</p>	<p>2</p> <p>and</p> <p>1</p>	<p>M1 for attempt to find 'gradient' using figures from the graph e.g. $(7.4 - 2.6) \div (2015 - 1951)$</p>	<p>Could be in billions Eg. $(7\,400\,000\,000 - 2\,600\,000\,000) \div (2015 - 1951)$</p> <p>For M1, condone incorrect conversion used consistently for both population figures.</p>
11	(a)		2	<p>B1 for $\frac{5}{9}$ and at least one fraction with denominator 8 for second card</p>	

Question		Answer	Marks	Part marks and guidance	
	(b)	$\frac{5}{9}$ oe	3	<p>M2FT for $\left(\frac{4}{9} \times \frac{5}{8}\right) + \left(\frac{5}{9} \times \frac{4}{8}\right)$ oe</p> <p>OR</p> <p>M1FT for $\left(\frac{4}{9} \times \frac{5}{8}\right)$ or $\left(\frac{5}{9} \times \frac{4}{8}\right)$ oe</p> <p>soi by $\frac{20}{72}$ oe</p>	FT <i>their</i> probabilities from (a)
12	(a)	5	1		
	(b)	(k =) 5 (r =) 1 nfw	5	<p>B1 for $206 = 41k + r$</p> <p>and</p> <p>B1 for $1031 = 206k + r$</p> <p>and</p> <p>M1 for $165k = 825$</p> <p>and</p> <p>A1 for $k = 5$ or $r = 1$</p> <p>If no or partial method shown, allow full marks for final answer correct</p>	<p>If 0 scored, allow SC2 for final correct answers interchanged</p> <p>Condone attempt to reduce to one variable by sub. or elim. With max of one error</p>
13	(a)	1.4355 or 1.436 or 1.44	2	M1 for 16.5×87 possibly soi by figs 14355, 1436 or 144	

Question		Answer	Marks	Part marks and guidance	
	(b)	Yes (Trevor is correct) because Eg $220 \div 87^3 \times 100^3 = 334.[\dots]$ or $334 \times 87^3 \div 100^3 = 219.9\dots$ to 220	3	M2 for $220 \div 87^3 \times 100^3$ or $334 \times 87^3 \div 100^3$ OR B1 for 87^3 or 658503 or 100^3 or 1000000 soi	
14	(a)	$(34 \times 36) - (25 \times 45) = 99$	2	M1 for either 34×36 or 25×45 soi by 1224 or 1125	

Question			Answer	Marks	Part marks and guidance
16	(a)	(i)	$\mathbf{b} - \mathbf{a}$	1	
		(ii)	$\frac{1}{4}(\mathbf{b} - \mathbf{a})$ or $\frac{1}{4}\mathbf{b} - \frac{1}{4}\mathbf{a}$	1	FT from (a)(i)
	(b)		$\overrightarrow{EF} = \overrightarrow{EB} + \overrightarrow{BF} = \frac{1}{4}(\mathbf{b} - \mathbf{a}) + \frac{1}{2}\mathbf{b}$ leading to $\frac{1}{4}(3\mathbf{b} - \mathbf{a})$ as given.	2	M1 for <i>their</i> part (a)(ii) + $\frac{1}{2}\mathbf{b}$ oe (a)(ii) must be in terms of a and b
	(c)		$\overrightarrow{AG} = \frac{3}{2}\mathbf{b} - \frac{1}{2}\mathbf{a}$ $\overrightarrow{AG} = 2\overrightarrow{EF}$ oe so are parallel.	3	B2 for $\overrightarrow{AG} = \frac{3}{2}\mathbf{b} - \frac{1}{2}\mathbf{a}$ or M1 for $\mathbf{b} + \frac{1}{2}$ (<i>their</i> part (a)(i)) oe Allow vectors found in reverse throughout eg. \overrightarrow{GA} instead of \overrightarrow{AG} Condone "AG and EF are multiples of each other" Full marks dependent on both AG and EF in correct simplified forms
17	(a)		$x^2 + y^2 = 100$ oe	1	
	(b)		$8^2 + (-6)^2 = 100$, so it's on the circle oe	2	M1 for $8^2 + (-6)^2$ seen or for substituting $x = 8$ and $y = -6$ into <i>their</i> part (a) <u>Alternative</u> using Pythagoras $\sqrt{8^2 + 6^2} = 10$ <i>their</i> part (a) must be an equation in both x and y.

Question		Answer	Marks	Part marks and guidance	
	(c)	$3y - 4x + 50 = 0$ oe	5	<p>B2 for [tangent gradient =] $\frac{4}{3}$ oe</p> <p>or</p> <p>M1 for $\pm \frac{6}{8}$ or $\pm \frac{8}{6}$ oe</p> <p>AND</p> <p>M2 for $y + 6 = \textit{their} \frac{4}{3}(x - 8)$ oe</p> <p>or</p> <p>M1 for $y = \textit{their} \frac{4}{3}x + 'c'$</p>	<p>Equivalents include:</p> $y = \frac{4}{3}x - \frac{50}{3}$ <p>Condone decimals with at least 2 decimal places rot: Eg. $y = 1.33x - 16.67$</p> <p>Equivalent for M2 includes $y = \textit{their} \frac{4}{3}x + c$ and then attempt to find c by substituting in $y = -6$ and $x = 8$</p>
18	(a)	$y \leq 2$ $y \geq -2x + 18$ oe	<p>1</p> <p>and</p> <p>3</p>	<p>B1 for ['gradient'=] -2 soi and</p> <p>M1 for suitable method to find equation of line eg. $y - 8 = (\textit{their} - 2) \times (x - 5)$ or $y - 2 = (\textit{their} - 2) \times (x - 8)$</p>	<p>If both inequalities are wrong way round, condone once (max penalty 1 mark)</p> <p>Or M1 for $y = \textit{their} -2x + c$ with a point from the line substituted in to find c For M1 allow use of an inequality symbol in place of =</p>
	(b)	$y = 6$ shown as a solid line and correct region shaded	2	<p>B1 for line drawn at $y = 6$</p> <p>OR</p> <p>B1 for correct squares shaded but no line</p>	Accept dashed line for B1

Question		Answer	Marks	Part marks and guidance	
	(c)	$\frac{8}{5}$ oe	5	<p>M1 for $\frac{1}{2} \times 4 \times (8 + 6)$ soi by 28</p> <p>M1 for $\frac{1}{2} \times 4h = \textit{their } 28 - 23$ oe</p> <p>A1 for $[h =] 2.5$</p> <p>AND</p> <p>M1 for $[k =] 4 \div \textit{their } 2.5$ oe</p> <p><u>Alternative method</u></p> <p>M1 for $\frac{1}{2} \times 4 \times (8 + 't')$</p> <p>M1 for <i>their</i> $\frac{1}{2} \times 4 \times (8 + 't') = 23$ oe</p> <p>A1 for $[t =] 3.5$</p> <p>AND</p> <p>M1 for $[k =] 4 \div (6 - \textit{their } 3.5)$ oe</p>	<p>'h' is 'top of triangle'</p> <p>'t' is 'top of trapezium'</p> <p>Must be a trapezium</p>

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2017

