



**1.** 
$$f(x) = \frac{1}{x(3x - 1)^2} = \frac{A}{x} + \frac{B}{(3x - 1)} + \frac{C}{(3x - 1)^2}$$

(a) Find the values of the constants  $A$ ,  $B$  and  $C$ .

**(4)**

(b) (i) Hence find  $\int f(x) \, dx$ .

(ii) Find  $\int_1^2 f(x) \, dx$ , leaving your answer in the form  $a + \ln b$ , where  $a$  and  $b$  are constants.

**(6)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









2.




Figure 1

Figure 1 shows a metal cube which is expanding uniformly as it is heated. At time  $t$  seconds, the length of each edge of the cube is  $x$  cm, and the volume of the cube is  $V$  cm<sup>3</sup>.

(a) Show that  $\frac{dV}{dx} = 3x^2$  (1)

Given that the volume,  $V$  cm<sup>3</sup>, increases at a constant rate of  $0.048$  cm<sup>3</sup>s<sup>-1</sup>,

(b) find  $\frac{dx}{dt}$ , when  $x = 8$  (2)

(c) find the rate of increase of the total surface area of the cube, in cm<sup>2</sup>s<sup>-1</sup>, when  $x = 8$  (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---











$$3. \quad f(x) = \frac{6}{\sqrt{(9-4x)}}, \quad |x| < \frac{9}{4}$$

- (a) Find the binomial expansion of  $f(x)$  in ascending powers of  $x$ , up to and including the term in  $x^3$ . Give each coefficient in its simplest form. (6)

Use your answer to part (a) to find the binomial expansion in ascending powers of  $x$ , up to and including the term in  $x^3$ , of

(b)  $g(x) = \frac{6}{\sqrt{(9+4x)}}, \quad |x| < \frac{9}{4}$  (1)

(c)  $h(x) = \frac{6}{\sqrt{(9-8x)}}, \quad |x| < \frac{9}{8}$  (2)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---











**Question 4 continued**

Lined writing area for the answer to Question 4.

**(Total 5 marks)**

**Q4**













6.




Figure 2

Figure 2 shows a sketch of the curve  $C$  with parametric equations

$$x = (\sqrt{3})\sin 2t, \quad y = 4 \cos^2 t, \quad 0 \leq t \leq \pi$$

(a) Show that  $\frac{dy}{dx} = k(\sqrt{3})\tan 2t$ , where  $k$  is a constant to be determined. (5)

(b) Find an equation of the tangent to  $C$  at the point where  $t = \frac{\pi}{3}$ .  
 Give your answer in the form  $y = ax + b$ , where  $a$  and  $b$  are constants. (4)

(c) Find a cartesian equation of  $C$ . (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







**Question 6 continued**

Lined area for student response. The area contains 25 horizontal lines for writing.

**(Total 12 marks)**

**Q6**

--	--



7.




Figure 3

Figure 3 shows a sketch of part of the curve with equation  $y = x^{\frac{1}{2}} \ln 2x$ .

The finite region  $R$ , shown shaded in Figure 3, is bounded by the curve, the  $x$ -axis and the lines  $x = 1$  and  $x = 4$

- (a) Use the trapezium rule, with 3 strips of equal width, to find an estimate for the area of  $R$ , giving your answer to 2 decimal places. (4)
- (b) Find  $\int x^{\frac{1}{2}} \ln 2x \, dx$ . (4)
- (c) Hence find the exact area of  $R$ , giving your answer in the form  $a \ln 2 + b$ , where  $a$  and  $b$  are exact constants. (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



















