13.



Figure 2

The resting heart rate, h, of a mammal, measured in beats per minute, is modelled by the equation

$$h = pm^q$$

where p and q are constants and m is the mass of the mammal measured in kg.

Figure 2 illustrates the linear relationship between $\log_{10} h$ and $\log_{10} m$

The line meets the vertical $\log_{10} h$ axis at 2.25 and has a gradient of -0.235

(a) Find, to 3 significant figures, the value of p and the value of q.

(3)

A particular mammal has a mass of 5 kg and a resting heart rate of 119 beats per minute.

(b) Comment on the suitability of the model for this mammal.

(3)

(c) With reference to the model, interpret the value of the constant p.

(1)

11. The owners of a nature reserve decided to increase the area of the reserve covered by trees.

Tree planting started on 1st January 2005.

The area of the nature reserve covered by trees, A km^2, is modelled by the equation

$$A = 80 - 45e^{ct}$$

where c is a constant and t is the number of years after 1st January 2005.

Using the model,

(a) find the area of the nature reserve that was covered by trees just before tree planting started.

(1)

On 1st January 2019 an area of 60 km^2 of the nature reserve was covered by trees.

(b) Use this information to find a complete equation for the model, giving your value of c to 3 significant figures.

(4)

On 1st January 2020, the owners of the nature reserve announced a long-term plan to have 100 km^2 of the nature reserve covered by trees.

(c) State a reason why the model is not appropriate for this plan.

(1)

5. A student's attempt to solve the equation $2\log_2 x - \log_2 \sqrt{x} = 3$ is shown below.

$2\log_2 x - \log_2 \sqrt{x} = 3$	
$2\log_2 \left(\frac{x}{\sqrt{x}}\right) = 3$	using the subtraction law for logs
$2\log_2 \left(\sqrt{x}\right) = 3$	simplifying
$\log_2 x = 3$	using the power law for logs
$x = 3^2 = 9$	using the definition of a log

(a) Identify two errors made by this student, giving a brief explanation of each.

(2)

(b) Write out the correct solution.

(3)

13.



Figure 3

The value of a rare painting, £V, is modelled by the equation $V = pq^t$, where p and q are constants and t is the number of years since the value of the painting was first recorded on 1st January 1980.

The line l shown in Figure 3 illustrates the linear relationship between t and $\log_{10} V$ since 1st January 1980.

The equation of line l is $\log_{10} V = 0.05t + 4.8$

(a) Find, to 4 significant figures, the value of p and the value of q.

(4)

(b) With reference to the model interpret

(i) the value of the constant p,

(ii) the value of the constant q.

(2)

(c) Find the value of the painting, as predicted by the model, on 1st January 2010, giving your answer to the nearest hundred thousand pounds.

(2)

14. The value of a car, £V, can be modelled by the equation

$$V = 15\,700e^{-0.25t} + 2300 \qquad t \in \mathbb{R},\ t \geqslant 0$$

where the age of the car is t years.

Using the model,

(a) find the initial value of the car.

(1)

Given the model predicts that the value of the car is decreasing at a rate of £500 per year at the instant when $t = T$,

(b) (i) show that

$$3925e^{-0.25T} = 500$$

(ii) Hence find the age of the car at this instant, giving your answer in years and months to the nearest month.

(*Solutions based entirely on graphical or numerical methods are not acceptable.*)

(6)

The model predicts that the value of the car approaches, but does not fall below, £A.

(c) State the value of A.

(1)

(d) State a limitation of this model.

(1)

8. The temperature, θ °C, of a cup of tea t minutes after it was placed on a table in a room, is modelled by the equation

$$\theta = 18 + 65e^{-\frac{t}{8}} \qquad t \geqslant 0$$

Find, according to the model,

(a) the temperature of the cup of tea when it was placed on the table,

(1)

(b) the value of t, to one decimal place, when the temperature of the cup of tea was 35 °C.

(3)

(c) Explain why, according to this model, the temperature of the cup of tea could not fall to 15 °C.

(1)



Figure 2

The temperature, μ °C, of a second cup of tea t minutes after it was placed on a table in a different room, is modelled by the equation

$$\mu = A + Be^{-\frac{t}{8}} \qquad t \geqslant 0$$

where A and B are constants.

Figure 2 shows a sketch of μ against t with two data points that lie on the curve.

The line l, also shown on Figure 2, is the asymptote to the curve.

Using the equation of this model and the information given in Figure 2

(d) find an equation for the asymptote l.

(4)

12. An advertising agency is monitoring the number of views of an online advert.

The equation

$$\log_{10} V = 0.072t + 2.379 \qquad 1 \leqslant t \leqslant 30,\ t \in \mathbb{N}$$

is used to model the total number of views of the advert, V, in the first t days after the advert went live.

(a) Show that $V = ab^t$ where a and b are constants to be found.

Give the value of a to the nearest whole number and give the value of b to 3 significant figures.

(4)

(b) Interpret, with reference to the model, the value of ab.

(1)

Using this model, calculate

(c) the total number of views of the advert in the first 20 days after the advert went live. Give your answer to 2 significant figures.

(2)

14.



Figure 2

A town's population, P, is modelled by the equation $P = ab^t$, where a and b are constants and t is the number of years since the population was first recorded.

The line l shown in Figure 2 illustrates the linear relationship between t and $\log_{10} P$ for the population over a period of 100 years.

The line l meets the vertical axis at (0, 5) as shown. The gradient of l is $\frac{1}{200}$.

(a) Write down an equation for l.

(2)

(b) Find the value of a and the value of b.

(4)

(c) With reference to the model interpret

(i) the value of the constant a,

(ii) the value of the constant b.

(2)

(d) Find

(i) the population predicted by the model when $t = 100$, giving your answer to the nearest hundred thousand,

(ii) the number of years it takes the population to reach 200 000, according to the model.

(3)

(e) State two reasons why this may not be a realistic population model.

(2)

Leave blank

7. (i) $2\log(x + a) = \log(16a^6)$, where a is a positive constant

Find x in terms of a, giving your answer in its simplest form.

(3)

(ii) $\log_3(9y + b) - \log_3(2y - b) = 2$, where b is a positive constant

Find y in terms of b, giving your answer in its simplest form.

(4)

Leave blank

8. (i) Given that

$$\log_3(3b + 1) - \log_3(a - 2) = -1, \qquad a > 2$$

express b in terms of a.

(3)

(ii) Solve the equation

$$2^{2x+5} - 7(2^x) = 0$$

giving your answer to 2 decimal places.
(*Solutions based entirely on graphical or numerical methods are not acceptable.*)

(4)

Leave blank

9. (i) Find the exact value of x for which

$$2\log_{10}(x-2) - \log_{10}(x+5) = 0$$

(5)

(ii) Given

$$\log_p(4y+1) - \log_p(2y-2) = 1 \qquad p > 2, y > 1$$

express y in terms of p.

(5)

Leave blank

5. (a) Given that

$$y = \log_3 x$$

find expressions in terms of y for

(i) $\log_3\left(\frac{x}{9}\right)$

(ii) $\log_3 \sqrt{x}$

Write each answer in its simplest form.

(3)

(b) Hence or otherwise solve

$$2\log_3\left(\frac{x}{9}\right) - \log_3 \sqrt{x} = 2$$

(4)

Leave blank

2. Find, giving your answer to 3 significant figures where appropriate, the value of x for which

(a) $7^{2x} = 14$

(3)

(b) $\log_5(3x + 1) = -2$

(2)

Leave blank

13. (a) Show that the equation

$$2\log_2 y = 5 - \log_2 x \qquad x > 0, y > 0$$

may be written in the form $y^2 = \frac{k}{x}$ where k is a constant to be found.

(3)

(b) Hence, or otherwise, solve the simultaneous equations

$$2\log_2 y = 5 - \log_2 x$$

$$\log_x y = -3$$

for $x > 0, y > 0$

(5)

Leave blank

14. (i) Given that

$$\log_a x + \log_a 3 = \log_a 27 - 1, \text{ where } a \text{ is a positive constant}$$

find, in its simplest form, an expression for x in terms of a.

(4)

(ii) Solve the equation

$$(\log_5 y)^2 - 7(\log_5 y) + 12 = 0$$

showing each step of your working.

(4)