




1.



**Figure 1**

A particle  $P$  of weight  $W$  newtons is attached to one end of a light inextensible string. The other end of the string is attached to a fixed point  $O$ . A horizontal force of magnitude 5 N is applied to  $P$ . The particle  $P$  is in equilibrium with the string taut and with  $OP$  making an angle of  $25^\circ$  to the downward vertical, as shown in Figure 1.

Find

- (a) the tension in the string, (3)  
(b) the value of  $W$ . (3)



## **Question 1 continued**

Leave  
blank



Leave  
blank

### **Question 1 continued**



### **Question 1 continued**

Leave  
blank

Q1

(Total 6 marks)



Leave  
blank

2. Two forces  $(4\mathbf{i} - 2\mathbf{j})$  N and  $(2\mathbf{i} + q\mathbf{j})$  N act on a particle  $P$  of mass 1.5 kg. The resultant of these two forces is parallel to the vector  $(2\mathbf{i} + \mathbf{j})$ .

- (a) Find the value of  $q$ .

(4)

At time  $t = 0$ ,  $P$  is moving with velocity  $(-2\mathbf{i} + 4\mathbf{j}) \text{ m s}^{-1}$ .

- (b) Find the speed of  $P$  at time  $t = 2$  seconds.

(6)



## **Question 2 continued**

Leave  
blank

Q2

(Total 10 marks)



3. A car starts from rest and moves with constant acceleration along a straight horizontal road. The car reaches a speed of  $V \text{ m s}^{-1}$  in 20 seconds. It moves at constant speed  $V \text{ m s}^{-1}$  for the next 30 seconds, then moves with constant deceleration  $\frac{1}{2} \text{ m s}^{-2}$  until it has speed  $8 \text{ m s}^{-1}$ . It moves at speed  $8 \text{ m s}^{-1}$  for the next 15 seconds and then moves with constant deceleration  $\frac{1}{3} \text{ m s}^{-2}$  until it comes to rest.

(a) Sketch, in the space below, a speed-time graph for this journey.

(3)

In the first 20 seconds of this journey the car travels 140 m.

Find

(b) the value of  $V$ ,

(2)

(c) the total time for this journey,

(4)

(d) the total distance travelled by the car.

(4)



### **Question 3 continued**

Leave  
blank



Leave  
blank

### **Question 3 continued**



### **Question 3 continued**

Leave  
blank

Q3

**(Total 13 marks)**



4. At time  $t = 0$ , a particle is projected vertically upwards with speed  $u$  from a point  $A$ . The particle moves freely under gravity. At time  $T$  the particle is at its maximum height  $H$  above  $A$ .

(a) Find  $T$  in terms of  $u$  and  $g$ .

(2)

(b) Show that  $H = \frac{u^2}{2g}$

(2)

The point  $A$  is at a height  $3H$  above the ground.

(c) Find, in terms of  $T$ , the total time from the instant of projection to the instant when the particle hits the ground.

(4)



### **Question 4 continued**

Leave  
blank



Leave  
blank

### **Question 4 continued**



## **Question 4 continued**


Leave  
blank

04

(Total 8 marks)



5.

**Figure 2**

Two particles  $A$  and  $B$  have masses  $2m$  and  $3m$  respectively. The particles are connected by a light inextensible string which passes over a smooth light fixed pulley. The system is held at rest with the string taut. The hanging parts of the string are vertical and  $A$  and  $B$  are above a horizontal plane, as shown in Figure 2. The system is released from rest.

- (a) Show that the tension in the string immediately after the particles are released is  $\frac{12}{5}mg$ . (6)

After descending 1.5 m,  $B$  strikes the plane and is immediately brought to rest. In the subsequent motion,  $A$  does not reach the pulley.

- (b) Find the distance travelled by  $A$  between the instant when  $B$  strikes the plane and the instant when the string next becomes taut. (6)

Given that  $m = 0.5 \text{ kg}$ ,

- (c) find the magnitude of the impulse on  $B$  due to the impact with the plane. (2)



## **Question 5 continued**

Leave  
blank



P 4 3 1 7 0 A 0 1 7 2 8

Leave  
blank

### **Question 5 continued**



## **Question 5 continued**


Leave  
blank



(Total 14 marks)



6.



**Figure 3**

A non-uniform beam  $AD$  has weight  $W$  newtons and length 4 m. It is held in equilibrium in a horizontal position by two vertical ropes attached to the beam. The ropes are attached to two points  $B$  and  $C$  on the beam, where  $AB = 1$  m and  $CD = 1$  m, as shown in Figure 3. The tension in the rope attached to  $C$  is double the tension in the rope attached to  $B$ . The beam is modelled as a rod and the ropes are modelled as light inextensible strings.

- (a) Find the distance of the centre of mass of the beam from  $A$ .

(6)

A small load of weight  $kW$  newtons is attached to the beam at  $D$ . The beam remains in equilibrium in a horizontal position. The load is modelled as a particle.

Find

- (b) an expression for the tension in the rope attached to  $B$ , giving your answer in terms of  $k$  and  $W$ , (3)

(c) the set of possible values of  $k$  for which both ropes remain taut. (2)



## Question 6 continued

Leave  
blank



Leave  
blank

## **Question 6 continued**



## **Question 6 continued**

Leave  
blank



(Total 11 marks)



7.




Figure 4

A particle  $P$  of mass  $2.7 \text{ kg}$  lies on a rough plane inclined at  $40^\circ$  to the horizontal. The particle is held in equilibrium by a force of magnitude  $15 \text{ N}$  acting at an angle of  $50^\circ$  to the plane, as shown in Figure 4. The force acts in a vertical plane containing a line of greatest slope of the plane. The particle is in equilibrium and is on the point of sliding down the plane.

Find

- (a) the magnitude of the normal reaction of the plane on  $P$ ,

(4)

- (b) the coefficient of friction between  $P$  and the plane.

(5)

The force of magnitude  $15 \text{ N}$  is removed.

- (c) Determine whether  $P$  moves, justifying your answer.

(4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



### **Question 7 continued**

Leave  
blank



Leave  
blank

### **Question 7 continued**



### **Question 7 continued**

Leave  
blank



Leave  
blank

### **Question 7 continued**

Q7

(Total 13 marks)

**TOTAL FOR PAPER: 75 MARKS**

END

