


# Mark Scheme (Results)

## June 2008

GCE

### GCE Mathematics (6683/01)


**June 2008  
6683 Statistics S1  
Mark Scheme**

Question Number	Scheme	Marks
Q1 (a)	 <p style="text-align: right; margin-right: 10%;">Tree without probabilities or labels 0.02(Disease), 0.95(Positive) on correct branches 0.03(Positive) on correct branch.</p>	<p><b>M1</b> <b>A1</b> <b>A1</b></p>
(b)	$P(\text{Positive Test}) = 0.02 \times 0.95 + 0.98 \times 0.03$ $= 0.0484$	<p><b>M1A1ft</b> <b>A1</b></p>
(c)	$P(\text{Do not have disease}   \text{Positive test}) = \frac{0.98 \times 0.03}{0.0484}$ $= 0.607438..$ <p style="text-align: right; margin-right: 10%;">awrt 0.607</p>	<p><b>[3]</b> <b>M1</b> <b>A1</b></p>
(d)	<p>Test not very useful OR High probability of not having the disease for a person with a positive test</p>	<p><b>[2]</b> <b>B1</b> <b>[1]</b></p>
<p><u>Notes:</u>                      (a) M1: All 6 branches.                      Bracketed probabilities not required.                      (b) M1 for sum of two products, at least one correct from their diagram                      A1ft follows from the probabilities on their tree                      A1 for correct answer only or <math>\frac{121}{2500}</math>                      (c) M1 for conditional probability with numerator following from their tree and denominator their answer to part (b).                      A1 also for <math>\frac{147}{242}</math>.</p>		<p><b>Total 9</b></p>

Question Number	Scheme	Marks
Q2 (a)	50	<b>B1</b>
(b)	$Q_1 = 45$ $Q_2 = 50.5$ $Q_3 = 63$	<b>B1</b> [1] <b>B1</b> <b>B1</b> <b>B1</b> [3]
(c)	$\text{Mean} = \frac{1469}{28} = 52.464286..$ $\text{Sd} = \sqrt{\frac{81213}{28} - \left(\frac{1469}{28}\right)^2}$ $= 12.164... \text{ or } 12.387216... \text{ for divisor } n-1$	ONLY awrt 52.5 <b>M1A1</b> awrt 12.2 or 12.4 <b>M1</b> <b>A1</b> [4]
(d)	$\frac{52.46.. - 50}{sd} = \text{awrt } 0.20 \text{ or } 0.21$	<b>M1A1</b> [2]
(e)	1. mode/median/mean Balmoral > mode/median/mean Abbey 2. Balmoral sd < Abbey sd or similar sd or correct comment from their values, Balmoral range < Abbey range, Balmoral IQR > Abbey IQR or similar IQR 3. Balmoral positive skew or almost symmetrical AND Abbey negative skew, Balmoral is less skew than Abbey or correct comment from their value in (d) 4. Balmoral residents generally older than Abbey residents or equivalent. Only one comment of each type max 3 marks	<b>B1B1B1</b> [3] <b>Total 13</b>
	<u>Notes:</u> (c) M1 for their 1469 between 1300 and 1600, divided by 28, A1 for awrt 52.5 .. Please note this is B1B1 on Epen M1 use of correct formula including sq root A1 awrt 12.2 or 12.4 Correct answers with no working award full marks. (d) M1 for their values correctly substituted A1 Accept 0.2 as a special case of awrt 0.20 with 0 missing (e) Technical terms required in correct context in lines 1 to 3 e.g. 'average' and 'spread' B0 1 correct comment B1B0B0 2 correct comments B1B1B0 3 correct comments B1B1B1	

Question Number	Scheme	Marks
<p>Q3</p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	$-1 \times p + 1 \times 0.2 + 2 \times 0.15 + 3 \times 0.15 = 0.55$ $p = 0.4$ $p + q + 0.2 + 0.15 + 0.15 = 1$ $q = 0.1$ $\text{Var}(X) = (-1)^2 \times p + 1^2 \times 0.2 + 2^2 \times 0.15 + 3^2 \times 0.15, -0.55^2$ $= 2.55 - 0.3025 = 2.2475$ $E(2X-4) = 2E(X) - 4$ $= -2.9$	<p><b>M1dM1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p>[5]</p> <p><b>M1A1,M1</b></p> <p><b>A1</b></p> <p>awrt 2.25</p> <p>[4]</p> <p><b>M1</b></p> <p><b>A1</b></p> <p>[2]</p> <p><b>Total 11</b></p>
	<p><u>Notes:</u></p> <p>(a) M1 for at least 2 correct terms on LHS                  Division by constant e.g. 5 then M0                  dM1 dependent on first M1 for equate to 0.55 and attempt to solve.                  Award M1M1A1 for <math>p=0.4</math> with no working                  M1 for adding probabilities and equating to 1. All terms or equivalent required                  e.g. <math>p+q=0.5</math>                  Award M1A1 for <math>q=0.1</math> with no working</p> <p>(b) M1 attempting <math>E(X^2)</math> with at least 2 correct terms                  A1 for fully correct expression or 2.55                  Division by constant at any point e.g. 5 then M0                  M1 for subtracting their mean squared                  A1 for awrt 2.25                  Award awrt 2.25 only with no working then 4 marks</p> <p>(c) M1 for <math>2x(\text{their mean}) - 4</math>                  Award 2 marks for -2.9 with no working</p>	

Question Number	Scheme	Marks
Q4 (a)	$S_{tt} = 10922.81 - \frac{401.3^2}{15} = 186.6973$	awrt 187 <b>M1A1</b>
	$S_{vv} = 42.3356 - \frac{25.08^2}{15} = 0.40184$	awrt 0.402 <b>A1</b>
	$S_{tv} = 677.971 - \frac{401.3 \times 25.08}{15} = 6.9974$	awrt 7.00 <b>A1</b>
(b)	$r = \frac{6.9974}{\sqrt{186.6973 \times 0.40184}} = 0.807869$	awrt 0.808 <b>M1A1ft</b> <b>A1</b>
(c)	<p><math>t</math> is the explanatory variable as we can control temperature but not frequency of noise or equivalent comment</p>	<b>B1</b> <b>B1</b>
(d)	<p>High value of <math>r</math> or <math>r</math> close to 1 or Strong correlation</p>	<b>B1</b>
(e)	$b = \frac{6.9974}{186.6973} = 0.03748$	awrt 0.0375 <b>M1A1</b>
	$a = \frac{25.08}{15} - b \times \frac{401.3}{15} = 0.6692874$	awrt 0.669 <b>M1A1</b>
(f)	$t = 19, v = 0.6692874 + 0.03748 \times 19 = 1.381406$	awrt 1.4 <b>B1</b>
	<p><u>Notes:</u></p> <p>(a) M1 any one attempt at a correct use of a formula. Award full marks for correct answers with no working. Epen order of awarding marks as above.</p> <p>(b) M1 for correct formula and attempt to use A1ft for their values from part (a)</p> <p>NB Special Case for <math>\frac{677.971}{\sqrt{10922.81 \times 42.3356}}</math> M1A0</p> <p>A1 awrt 0.808 Award 3 marks for awrt 0.808 with no working</p> <p>(c) Marks are independent. Second mark requires some interpretation in context and can be statements such as ‘temperature effects / influences pitch or noise’ B1 ‘temperature is being changed’ BUT B0 for ‘temperature is changing’</p> <p>(e) M1 their values the right way up A1 for awrt 0.0375 M1 attempt to use correct formula with their value of <math>b</math> A1 awrt 0.669</p> <p>(f) awrt 1.4</p>	<b>[4]</b> <b>[3]</b> <b>[2]</b> <b>[1]</b> <b>[4]</b> <b>[1]</b> <b>Total 15</b>

Question Number	Scheme	Marks
<p>Q5 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<div style="display: flex; align-items: center; justify-content: center;">  <div style="margin-left: 20px;"> <p>3 closed intersecting curves with labels 100 100,30 12,10,3,25 Box</p> </div> </div> <p>(b) <math>P(\text{Substance } C) = \frac{100+100+10+25}{300} = \frac{235}{300} = \frac{47}{60}</math> or exact equivalent</p> <p>(c) <math>P(\text{All 3}   A) = \frac{10}{30+3+10+100} = \frac{10}{143}</math> or exact equivalent</p> <p>(d) <math>P(\text{Universal donor}) = \frac{20}{300} = \frac{1}{15}</math> or exact equivalent</p>	<p><b>M1</b> <b>A1</b> <b>A1</b> <b>B1</b></p> <p style="text-align: right;"><b>[4]</b></p> <p><b>M1A1ft</b> <b>[2]</b></p> <p><b>M1A1ft</b> <b>[2]</b></p> <p><b>M1A1 cao</b> <b>[2]</b></p> <p><b>Total 10</b></p>
	<p><u>Notes:</u></p> <p>(a) 20 not required. Fractions and exact equivalent decimals or percentages.</p> <p>(b) M1 For adding their positive values in C and finding a probability A1ft for correct answer or answer from their working</p> <p>(c) M1 their 10 divided by their sum of values in A A1ft for correct answer or answer from their working</p> <p>(d) M1 for 'their 20' divided by 300 A1 correct answer only</p>	

Question Number	Scheme	Marks								
<p>Q6 (a)</p> <p>(b)</p>	<p>F(4)=1  <math>(4+k)^2 = 25</math>  <math>k = 1</math> as <math>k &gt; 0</math></p> <table border="1" data-bbox="220 472 1326 595"> <tr> <td style="text-align: center;"><math>x</math></td> <td style="text-align: center;">2</td> <td style="text-align: center;">3</td> <td style="text-align: center;">4</td> </tr> <tr> <td style="text-align: center;"><math>P(X=x)</math></td> <td style="text-align: center;"><math>\frac{9}{25}</math></td> <td style="text-align: center;"><math>\frac{7}{25}</math></td> <td style="text-align: center;"><math>\frac{9}{25}</math></td> </tr> </table>	$x$	2	3	4	$P(X=x)$	$\frac{9}{25}$	$\frac{7}{25}$	$\frac{9}{25}$	<p><b>M1</b>  <b>A1</b></p> <p style="text-align: right;"><b>[2]</b></p> <p><b>B1ftB1B1</b></p> <p style="text-align: right;"><b>[3]</b></p> <p><b>Total 5</b></p>
$x$	2	3	4							
$P(X=x)$	$\frac{9}{25}$	$\frac{7}{25}$	$\frac{9}{25}$							
	<p><u>Notes:</u>                      (a) M1 for use of <math>F(4) = 1</math> only If <math>F(2)=1</math> and / or <math>F(3)=1</math> seen then M0.  <math>F(2)+F(3)+F(4)=1</math> M0                      A1 for <math>k=1</math> and ignore <math>k= -9</math></p> <p>(b) B1ft follow through their <math>k</math> for <math>P(X=2)</math> either exact or 3sf between 0 and 1 inclusive.                      B1 correct answer only or exact equivalent                      B1 correct answer only or exact equivalent</p>									

Question Number	Scheme	Marks
<p>Q7 (a)</p> <p>(b)</p> <p>(c)</p>	$z = \frac{53 - 50}{2}$ $P(X > 53) = 1 - P(Z < 1.5)$ $= 1 - 0.9332$ $= 0.0668$ $P(X \leq x_0) = 0.01$ $\frac{x_0 - 50}{2} = -2.3263$ $x_0 = 45.3474$ $P(2 \text{ weigh more than } 53\text{kg and } 1 \text{ less}) = 3 \times 0.0668^2 (1 - 0.0668)$ $= 0.012492487..$	<p>Attempt to standardise 1-probability required can be implied</p> <p><b>M1</b> <b>B1</b> <b>A1</b></p> <p><b>M1</b></p> <p><b>M1B1</b> <b>M1A1</b></p> <p><b>B1M1A1ft</b> <b>A1</b></p> <p><b>Total 12</b></p> <p>[3] [5] [4]</p>
	<p><u>Notes:</u>                      (a) M1 for using 53,50 and 2, either way around on numerator                      B1 1- any probability for mark                      A1 0.0668 cao                      (b) M1 can be implied or seen in a diagram                      or equivalent with correct use of 0.01 or 0.99                      M1 for attempt to standardise with 50 and 2 numerator either way around                      B1 for <math>\pm 2.3263</math>                      M1 Equate expression with 50 and 2 to a z value to form an equation with consistent signs and attempt to solve                      A1 awrt 45.3 or 45.4                      (c) B1 for 3,                      M1 <math>p^2(1 - p)</math> for any value of <math>p</math>                      A1ft for <math>p</math> is their answer to part (a) without 3                      A1 awrt 0.012 or 0.0125</p>	