

# Mark Scheme (Results)

Summer 2012

GCE Core Mathematics C2  
(6664) Paper 1

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2012

Publications Code UA031953

All the material in this publication is copyright

© Pearson Education Ltd 2012

**Summer 2012**  
**6664 Core Mathematics**  
**C2 Mark Scheme**

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## EDEXCEL GCE MATHEMATICS

### General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\checkmark$  will be used for correct ft
  - cao – correct answer only
  - cso – correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - $\square$  The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

## General Principles for Core Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

### Method mark for solving 3 term quadratic:

#### 1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$ , leading to  $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$ , leading to  $x = \dots$

#### 2. Formula

Attempt to use correct formula (with values for  $a$ ,  $b$  and  $c$ ), leading to  $x = \dots$

#### 3. Completing the square

Solving  $x^2 + bx + c = 0$ :  $(x \pm \frac{b}{2})^2 \pm q \pm c$ ,  $q \neq 0$ , leading to  $x = \dots$

### Method marks for differentiation and integration:

#### 1. Differentiation

Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )

#### 2. Integration

Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

### Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

### Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

### Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required.

## Summer 2012 6664 Core Mathematics 2 Mark Scheme

Question number	Scheme	Marks
<b>1</b>	$[(2-3x)^5] = \dots + \binom{5}{1} 2^4 (-3x) + \binom{5}{2} 2^3 (-3x)^2 + \dots, \dots$ $= 32, -240x, +720x^2$	<p>M1</p> <p>B1, A1, A1</p>
Notes	<b>Total 4</b>	
	<p><b>M1:</b> The <b>method</b> mark is awarded for an attempt at Binomial to get the second and/or third term – need <b>correct</b> binomial coefficient combined <b>with correct power of x</b>. Ignore errors (or omissions) in powers of 2 or 3 or sign or bracket errors. Accept any notation for <math>{}^5C_1</math> and <math>{}^5C_2</math>, e.g. <math>\binom{5}{1}</math> and <math>\binom{5}{2}</math> (unsimplified) or 5 and 10 from Pascal's triangle This mark may be given if no working is shown, but either or both of the terms including <math>x</math> is correct.</p> <p><b>B1:</b> must be simplified to 32 ( writing just <math>2^5</math> is <b>B0</b> ). <b>32</b> must be the only constant term in the final answer- so <math>32 + 80 - 3x + 80 + 9x^2</math> is B0 but may be eligible for M1A0A0 .</p> <p><b>A1:</b> is <b>cao</b> and is for <math>-240x</math>. (not <math>+240x</math>) The <math>x</math> is required for this mark</p> <p><b>A1:</b> is c.a.o and is for <math>720x^2</math> (can follow omission of negative sign in working)</p> <p>A list of correct terms may be given credit i.e. series appearing on different lines</p> <p>Ignore extra terms in <math>x^3</math> and/or <math>x^4</math> (isw)</p>	
Special Case	<p>Special Case: <i>Descending powers</i> of <math>x</math> would be</p> <p><math>(-3x)^5 + 2 \times 5 \times (-3x)^4 + 2^2 \times \binom{5}{3} \times (-3x)^3 + \dots</math> i.e. <math>-243x^5 + 810x^4 - 1080x^3 + \dots</math> This is a misread but award as s.c. M1B1A0A0 if completely “correct” or M1 B0A0A0 for <u>correct binomial coefficient</u> in any form with the correct power of <math>x</math></p>	
Alternative Method	<p><b>Method 1:</b> <math>[(2-3x)^5] = 2^5 \left( 1 + \binom{5}{1} \left(-\frac{3x}{2}\right) + \binom{5}{2} \left(\frac{-3x}{2}\right)^2 + \dots \right)</math> is <b>M1B0A0A0</b> { The M1 is for the expression <b>in the bracket</b> and as in first method– need <b>correct</b> binomial coefficient combined with correct power of <math>x</math>. Ignore bracket errors or errors (or omissions) in powers of 2 or 3 or sign or bracket errors }</p> <p>– answers must be simplified to <math>= 32, -240x, +720x^2</math> for full marks (awarded as before)</p> <p><math>[(2-3x)^5] = 2 \left( 1 + \binom{5}{1} \left(-\frac{3x}{2}\right) + \binom{5}{2} \left(\frac{-3x}{2}\right)^2 + \dots \right)</math> would also be awarded <b>M1B0A0A0</b></p> <p><b>Method 2: Multiplying out :</b> B1 for 32 and M1A1A1 for other terms with M1 awarded if <math>x</math> or <math>x^2</math> term is correct. Completely correct is 4/4</p>	

Question number	Scheme	Marks
2	$2 \log x = \log x^2$ $\log_3 x^2 - \log_3(x-2) = \log_3 \frac{x^2}{x-2}$ $\frac{x^2}{x-2} = 9$ <p>Solves <math>x^2 - 9x + 18 = 0</math> to give <math>x = \dots</math></p> $x = 3, x = 6$	B1 M1 A1 o.e. M1 A1 <hr/> <b>Total 5</b>
Notes	<p><b>B1</b> for <b>this</b> correct use of power rule (may be implied)  <b>M1</b>: for <b>correct</b> use of subtraction rule (or addition rule) for logs            N.B. <math>2 \log_3 x - \log_3(x-2) = 2 \log_3 \frac{x}{x-2}</math> is <b>M0</b>  <b>A1</b>. for correct equation without logs (Allow any correct equivalent including <math>3^2</math> instead of 9.)  <b>M1</b> for attempting to solve <math>x^2 - 9x + 18 = 0</math> to give <math>x =</math> (see notes on marking quadratics)  <b>A1</b> for these <b>two</b> correct answers</p>	
Alternative Method	<p><math>\log_3 x^2 = 2 + \log_3(x-2)</math> is B1,  <i>so</i> <math>x^2 = 3^{2+\log_3(x-2)}</math> needs to be followed by <math>(x^2) = 9(x-2)</math> for M1 A1            Here M1 is for <b>complete</b> method i.e. correct use of powers after logs are used correctly</p>	
Common Slips	<p><math>2 \log x - \log x + \log 2 = 2</math> may obtain B1 if <math>\log x^2</math> appears but <b>the statement is M0</b> and so leads to no further marks  <math>2 \log_3 x - \log_3(x-2) = 2</math> <i>so</i> <math>\log_3 x - \log_3(x-2) = 1</math> and <math>\log_3 \frac{x}{x-2} = 1</math> can earn M1 for <i>correct</i> subtraction rule following error, but no other marks</p>	
Special Case	<p><math>\frac{\log x^2}{\log(x-2)} = 2</math> leading to <math>\frac{x^2}{x-2} = 9</math> and then to <math>x=3, x=6</math>, usually earns B1M0A0, but may then earn M1A1 (special case) so 3/5 [ This <i>recovery</i> after uncorrected error is very common]            Trial and error, Use of a table or just stating answer <b>with both</b> <math>x=3</math> and <math>x=6</math> should be awarded B0M0A0 then final M1A1 i.e. 2/5</p>	

Question number	Scheme	Marks
3	Obtain $\underline{(x \pm 10)^2}$ <b>and</b> $\underline{(y \pm 8)^2}$	M1
(a)	Obtain $\underline{(x - 10)^2}$ <b>and</b> $\underline{(y - 8)^2}$	A1
	Centre is (10, 8). N.B. This may be indicated on <b>diagram only</b> as (10, 8)	A1
(b)	See $\underline{(x \pm 10)^2} + \underline{(y \pm 8)^2} = 25 (= r^2)$ or $(r^2 =) "100" + "64" - 139$ $r = 5$ * (this is a printed answer so need one of the above two reasons)	M1 A1
(c)	Use $x = 13$ in either form of equation of circle and solve resulting quadratic to give $y =$ e.g. $x = 13 \Rightarrow (13 - 10)^2 + (y - 8)^2 = 25 \Rightarrow (y - 8)^2 = 16$ so $y =$ or $13^2 + y^2 - 20 \times 13 - 16y + 139 = 0 \Rightarrow y^2 - 16y + 48 = 0$ so $y =$ $y = 4$ or $12$ ( on EPEN mark one correct value as A1A0 and both correct as A1 A1)	(3) M1 A1
(d)	Use of $r\theta$ with $r = 5$ and $\theta = 1.855$ (may be implied by 9.275)  Perimeter $PTQ = 2r +$ their <b>arc</b> $PQ$ (Finding perimeter of triangle is M0 here)  $= 19.275$ or $19.28$ or $19.3$	M1 A1 A1 (3)
		<b>11 marks</b>
<b>Alternatives</b>	<i>Method 2:</i> From $x^2 + y^2 + 2gx + 2fy + c = 0$ centre is $(\pm g, \pm f)$	M1
(a)	Centre is $(-g, -f)$ , <b>and so centre is</b> (10, 8).	A1, A1
OR	<i>Method 3:</i> Use any value of $y$ to give two points ( $L$ and $M$ ) on circle. $x$ co-ordinate of mid point of $LM$ is "10" <b>and</b> Use any value of $x$ to give two points ( $P$ and $Q$ ) on circle. $y$ co-ordinate of mid point of $PQ$ is "8" (Centre – chord theorem) . (10,8) is M1A1A1	M1 A1 A1 (3)
(b)	<i>Method 2:</i> Using $\sqrt{g^2 + f^2 - c}$ or $(r^2 =) "100" + "64" - 139$ $r = 5$ *	M1 A1
OR	<i>Method 3:</i> Use point on circle with centre to find radius. Eg $\sqrt{(13 - 10)^2 + (12 - 8)^2}$ $r = 5$ *	M1 A1 cao (2)
(c)	Divide triangle $PTQ$ and use Pythagoras with $r^2 - (13 - "10")^2 = h^2$ , then evaluate "8 $\pm$ h" - (N.B. Could use 3,4,5 Triangle and $8 \pm 4$ ). Accuracy as before	M1
<b>Notes</b>	<b>Mark (a) and (b) together</b>	
(a)	<b>M1</b> as in scheme and can be <u>implied</u> by $(\pm 10, \pm 8)$ . <b>Correct centre (10, 8) implies M1A1A1</b>	
(b)	<b>M1</b> for a correct method leading to $r = \dots$ , or $r^2 = "100" + "64" - 139$ (not $139 - "100" - "64"$ ) or for using equation of circle in $\underline{(x \pm 10)^2} + \underline{(y \pm 8)^2} = k^2$ form to identify $r =$ <b>3<sup>rd</sup> A1</b> $r = 5$ ( <b>NB This is a given answer so should follow</b> $k^2 = 25$ or $r^2 = 100 + 64 - 139$ ) <b>Special case:</b> if centre is given as (-10, -8) or (10, -8) or (-10, 8) allow <b>M1A1</b> for $r = 5$ worked correctly as $r^2 = 100 + 64 - 139$	
(d)	Full marks available for calculation using major sector so Use of $r\theta$ with $r = 5$ and $\theta = 4.428$ leading to perimeter of 32.14 for major sector	



Question number	Scheme	Marks
<p><b>4 (a)</b></p> <p><b>(b)</b></p>	$f(-2) = 2 \cdot (-2)^3 - 7 \cdot (-2)^2 - 10 \cdot (-2) + 24$ $= 0 \text{ so } (x+2) \text{ is a factor}$ $f(x) = (x+2)(2x^2 - 11x + 12)$ $f(x) = (x+2)(2x-3)(x-4)$	<p>M1 A1 (2)</p> <p>M1 A1 dM1 A1 (4)</p> <p><b>6 marks</b></p>
<p>Notes (a)</p> <p><b>(b)</b></p>	<p><b>M1</b> : Attempts <math>f(\pm 2)</math> (Long division is M0)  <b>A1</b> : is for <math>=0</math> <b>and conclusion</b>  <b>Note</b>: Stating “hence factor” or “it is a factor” or a “<math>\surd</math>” (tick) or “QED” is fine for the conclusion.  <b>Note</b> also that a conclusion can be implied from a <u>preamble</u>, eg: “If <math>f(-2) = 0</math>, <math>(x + 2)</math> is a factor...” (Not just <math>f(-2)=0</math>)</p> <p><b>1<sup>st</sup> M1</b>: Attempts long division by correct factor or other method leading to obtaining <math>(2x^2 \pm ax \pm b)</math>, <math>a \neq 0</math>, <math>b \neq 0</math>, even with a remainder. Working need not be seen as could be done “by inspection.”</p> <p>Or <i>Alternative Method</i> : <b>1<sup>st</sup> M1</b>: Use <math>(x+2)(ax^2 + bx + c) = 2x^3 - 7x^2 - 10x + 24</math> with expansion and comparison of coefficients to obtain <math>a = 2</math> and to obtain values for <math>b</math> and <math>c</math></p> <p><b>1<sup>st</sup> A1</b>: For seeing <math>(2x^2 - 11x + 12)</math>. [Can be seen here in (b) after work done in (a)]</p> <p><b>2<sup>nd</sup> M1</b>: Factorises quadratic. (see rule for factorising a quadratic). This is dependent on the previous method mark being awarded and <b>needs factors</b></p> <p><b>2<sup>nd</sup> A1</b>: is cao and needs all three factors together. Ignore subsequent work (such as a solution to a quadratic equation.)</p> <p>Note: Some candidates will go from <math>\{(x+2)\}(2x^2 - 11x + 12)</math> to <math>\{x = -2\}</math>, <math>x = \frac{3}{2}</math>, <math>4</math>, and not list all three factors. Award these responses M1A1M0A0.</p> <p>Finds <math>x = 4</math> and <math>x = 1.5</math> by factor theorem, formula or calculator and produces factors M1  <math>f(x) = (x+2)(2x-3)(x-4)</math> <b>or</b> <math>f(x) = 2(x+2)(x-1.5)(x-4)</math> <b>o.e. is full marks</b>  <math>f(x) = (x+2)(x-1.5)(x-4)</math> <b>loses last A1</b></p>	

Question number	Scheme		Marks
<p>Method 1 5 (a)</p> <p>(b)</p>	<p>Puts <math>10 - x = 10x - x^2 - 8</math> and rearranges to give three term quadratic</p> <p>Solves their "<math>x^2 - 11x + 18 = 0</math>" using acceptable method as in general principles to give <math>x =</math></p> <p>Obtains <math>x = 2, x = 9</math> (may be on diagram or in part (b) in limits)</p> <p>Substitutes their <math>x</math> into a given equation to give <math>y =</math> (may be on diagram)</p> <p><math>y = 8, y = 1</math></p> $\int (10x - x^2 - 8) dx = \frac{10x^2}{2} - \frac{x^3}{3} - 8x \{+ c\}$ $\left[ \frac{10x^2}{2} - \frac{x^3}{3} - 8x \right]_2^9 = (\dots) - (\dots)$ $= 90 - \frac{4}{3} = 88\frac{2}{3} \text{ or } \frac{266}{3}$ <p>Area of trapezium = <math>\frac{1}{2}(8+1)(9-2) = 31.5</math></p> <p>So area of <math>R</math> is <math>88\frac{2}{3} - 31.5 = 57\frac{1}{6}</math> or <math>\frac{343}{6}</math></p>	<p>Or puts <math>y = 10(10 - y) - (10 - y)^2 - 8</math> and rearranges to give three term quadratic</p> <p>Solves their "<math>y^2 - 9y + 8 = 0</math>" using acceptable method as in general principles to give <math>y =</math></p> <p>Obtains <math>y = 8, y = 1</math> (may be on diagram)</p> <p>Substitutes their <math>y</math> into a given equation to give <math>x =</math> (may be on diagram or in part (b))</p> <p><math>x = 2, x = 9</math></p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 (5)</p> <p>M1 A1 A1</p> <p>dM1</p> <p>B1</p> <p>M1A1 cao (7)</p> <p><b>12 marks</b></p>
<p><b>Notes</b> (a)</p> <p>(b)</p>	<p>First <b>M1</b>: See scheme Second <b>M1</b>: See notes relating to solving quadratics</p> <p>Third <b>M1</b>: This may be awarded if one substitution is made</p> <p>Two correct Answers following tables of values, or from Graphical calculator are 5/5</p> <p><b>Just one pair of correct coordinates – no working or from table is M0M0A0M1A0</b></p> <p><b>M1</b>: <math>x^n \rightarrow x^{n+1}</math> for any one term.</p> <p><b>1<sup>st</sup> A1</b>: at least two out of three terms correct <b>2<sup>nd</sup> A1</b>: All three correct</p> <p><b>dM1</b>: Substitutes 9 and 2 (or limits from part(a)) into an "integrated function" and subtracts, either way round</p> <p>(NB: If candidate <b>changes all signs</b> to get <math>\int (-10x + x^2 + 8) dx = -\frac{10x^2}{2} + \frac{x^3}{3} + 8x \{+ c\}</math> This is M1 A1 A1</p> <p>Then uses limits dM1 and trapezium is B1</p> <p>Needs to <i>change sign of value obtained</i> from integration for final M1A1 so <math>-88\frac{2}{3} - 31.5</math> is M0A0 )</p> <p><b>B1</b>: Obtains 31.5 for area under line using any correct method (could be integration) or triangle minus triangle <math>\frac{1}{2} \times 8 \times 8 - \frac{1}{2}</math> or rectangle plus triangle [may be implied by correct 57 1/6 ]</p> <p><b>M1</b>: Their Area under curve – Their Area under line (if integrate both need same limits)</p> <p><b>A1</b>: Accept 57.16recurring but not 57.16</p> <p><b>PTO for Alternative method</b></p>		

<p><b>Method 2 for (b)</b></p>	<p>Area of R</p> $= \int_2^9 (10x - x^2 - 8) - (10 - x) dx$ $\int_2^9 -x^2 + 11x - 18 dx$ $= -\frac{x^3}{3} + \frac{11x^2}{2} - 18x \{+ c\}$ $\left[ -\frac{x^3}{3} + \frac{11x^2}{2} - 18x \right]_2^9 = (\dots) - (\dots)$ <p>This mark is implied by final answer which rounds to 57.2</p> <p><i>See above working(allow bracketing errors) to decide to award 3<sup>rd</sup> M1 mark for (b) here:</i></p> $40.5 - (-16\frac{2}{3}) = 57\frac{1}{6} \text{ cao}$	<p>3<sup>rd</sup> M1 (in (b) ): Uses difference between two functions in integral.  M: <math>x^n \rightarrow x^{n+1}</math> for any one term.  A1 at least two out of these three simplified terms  Correct integration. (Ignore + c).  Substitutes 9 and 2 (or limits from part(a)) into an "integrated function" and subtracts, either way round.</p>	<p>M1  A1  A1  dM1  B1  M1  A1  (7)</p>
<p>Special case of above method</p>	$\int_2^9 x^2 - 11x + 18 dx = \frac{x^3}{3} - \frac{11x^2}{2} + 18x \{+ c\}$ $\left[ \frac{x^3}{3} - \frac{11x^2}{2} + 18x \right]_2^9 = (\dots) - (\dots)$ <p>This mark is implied by final answer which rounds to 57.2 (not -57.2)</p> <p>Difference of functions implied (see above expression)</p> $40.5 - (-16\frac{2}{3}) = 57\frac{1}{6} \text{ cao}$		<p>M1A1A1  DM1  B1  M1  A1  (7)</p>
<p>Special Case 2</p>	<p>Integrates expression in y e.g. "<math>y^2 - 9y + 8 = 0</math>": This can have first M1 in part (b) and no other marks. (It is not a method for finding this area)</p>		
<p>Notes</p>	<p>Take away trapezium again having used Method 2 loses last two marks</p> <p>Common Error:</p> <p>Integrates <math>-x^2 + 9x - 18</math> is likely to be M1A1A0dM1B0M1A0</p> <p>Integrates <math>2 - 11x - x^2</math> is likely to be M1A0A0dM1B0M1A0</p> <p>Writing <math>\int_2^9 (10x - x^2 - 8) - (10 - x) dx</math> only earns final M mark</p>		

Question number	Scheme	Marks
6(a)	States or uses $\tan 2x = \frac{\sin 2x}{\cos 2x}$  $\frac{\sin 2x}{\cos 2x} = 5 \sin 2x \Rightarrow \sin 2x - 5 \sin 2x \cos 2x = 0 \Rightarrow \sin 2x(1 - 5 \cos 2x) = 0$ *	M1  A1  (2)
(b)	$\sin 2x = 0$ gives $2x = 0, 180, 360$ so $x = 0, 90, 180$  $\cos 2x = \frac{1}{5}$ gives $2x = 78.46$ (or 78.5 or 78.4) <b>or</b> $2x = 281.54$ (or 281.6)  $x = 39.2$ (or 39.3), $140.8$ (or 141)	B1 for two correct answers, second B1 for all three correct. Excess in range – lose last B1  B1, B1  M1  A1, A1  (5)  <b>7 marks</b>
Notes	<p>(a) <b>M1</b>: Statement that <math>\tan \theta = \frac{\sin \theta}{\cos \theta}</math> or Replacement of tan (wherever it appears). Must be a correct statement but may involve <math>\theta</math> instead of <math>2x</math>.  <b>A1</b>: the answer is given so all steps should be given.</p> <p>N.B. <math>\sin 2x - 5 \sin 2x \cos 2x = 0</math> or <math>-5 \sin 2x \cos 2x + \sin 2x = 0</math> or <math>\sin 2x(\frac{1}{\cos 2x} - 5) = 0</math> o.e.</p> <p><b>must be seen</b> and be followed by printed answer for A1 mark  <math>\sin 2x = 5 \sin 2x \cos 2x</math> is not sufficient.</p> <p>(b) Statement of 0 and 180 with no working gets B1 B0 (bod) as it is two solutions  <b>M1</b>: This mark for one of the two statements given (must relate to <math>2x</math> not just to <math>x</math>)  <b>A1, A1</b>: first A1 for 39.2, second for 140.8  <i>Special case</i> solving <math>\cos 2x = -1/5</math> giving <math>2x = 101.5</math> or <math>258.5</math> is awarded M1A0A0                      140.8 omitted would give M1A1A0                      Allow answers which round to 39.2 or 39.3 and which round to 140.8 and allow 141                      Answers in radians lose last A1 awarded (These are 0, 0.68, 1.57, 2.46 and 3.14)                      Excess answers <b>in range</b> lose last A1 Ignore excess answers outside range.                      All 5 correct answers with no extras and no working gets <b>full marks</b> in part (b). The answers imply the method here</p>	

Question number	Scheme	Marks												
7 (a)	<table border="1" data-bbox="312 387 1010 465"> <tr> <td><math>x</math></td> <td>0</td> <td>0.25</td> <td>0.5</td> <td>0.75</td> <td>1</td> </tr> <tr> <td><math>y</math></td> <td>1</td> <td>1.251</td> <td><b>1.494</b></td> <td><b>1.741</b></td> <td>2</td> </tr> </table>	$x$	0	0.25	0.5	0.75	1	$y$	1	1.251	<b>1.494</b>	<b>1.741</b>	2	B1, B1 (2)
$x$	0	0.25	0.5	0.75	1									
$y$	1	1.251	<b>1.494</b>	<b>1.741</b>	2									
(b)	$\frac{1}{2} \times 0.25, \{(1+2) + 2(1.251+1.494+1.741)\} \text{ o.e.}$ $=1.4965$	B1, M1,A1 ft  A1 (4)												
		<b>6 marks</b>												
Notes	<p>(a) first <b>B1</b> for 1.494 and second <b>B1</b> for 1.741 ( 1.740 is <b>B0</b> ) Wrong accuracy e.g. 1.49, 1.74 is B1B0</p> <p>(b) <b>B1</b>: Need <math>\frac{1}{2}</math> of 0.25 or 0.125 o.e. <b>M1</b>: requires first bracket to contain first plus last values <b>and</b> second bracket to include no additional values from the three in the table. If the only mistake is to omit one value from second bracket this may be regarded as a slip and <b>M</b> mark can be allowed ( An extra repeated term forfeits the <b>M</b> mark however) <math>x</math> values: <b>M0</b> if values used in brackets are <math>x</math> values instead of <math>y</math> values</p> <p><b>A1ft</b> follows their answers to part (a) and is for {correct expression} Final <b>A1</b>: Accept 1.4965, 1.497. or 1.50 only after correct work. (No follow through except one special case below following 1.740 in table) Separate trapezia may be used : <b>B1</b> for 0.125, <b>M1</b> for <math>\frac{1}{2}h(a+b)</math> used 3 or 4 times (and <b>A1ft</b> if it is all correct ) e.g.. <math>0.125(1+ 1.251) + 0.125(1.251+1.494) + 0.125(1.741 + 2)</math> is <b>M1 A0</b> equivalent to missing one term in { } in main scheme</p> <p>Special Case: Bracketing mistake: i.e. <math>0.125(1+2) + 2(1.251+1.494+1.741)</math> scores <b>B1 M1 A0 A0 for 9.347</b> If the final answer implies that the calculation has been done correctly i.e. 1.4965 (then full marks can be given). <b>Need to see trapezium rule – answer only (with no working) is 0/4 any doubts send to review</b></p> <p>Special Case; Uses 1.740 to give 1.49625 or 1.4963 or 1.496 or 1.50 gets, B1 B0 B1M1A1ft then A1 (lose 1 mark)</p> <p>NB Bracket is 11.972</p>													

Question number	Scheme	Marks
<p>8</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p><math>(h =) \frac{60}{\pi x^2}</math> or <b>equivalent exact</b> (not decimal) expression e.g. <math>(h =) 60 \div \pi x^2</math></p> <p><math>(A =) 2\pi x^2 + 2\pi xh</math> or <math>(A =) 2\pi r^2 + 2\pi rh</math> or <math>(A =) 2\pi r^2 + \pi dh</math> may not be simplified and may appear on separate lines</p> <p>Either <math>(A) = 2\pi x^2 + 2\pi x \left( \frac{60}{\pi x^2} \right)</math> or As <math>\pi xh = \frac{60}{x}</math> then <math>(A =) 2\pi x^2 + 2 \left( \frac{60}{x} \right)</math></p> <p><math>A = 2\pi x^2 + \left( \frac{120}{x} \right)</math> *</p> <p><math>\left( \frac{dA}{dx} \right) = 4\pi x - \frac{120}{x^2}</math> or <math>= 4\pi x - 120x^{-2}</math></p> <p><math>4\pi x - \frac{120}{x^2} = 0</math> implies <math>x^3 =</math> (Use of <math>&gt; 0</math> or <math>&lt; 0</math> is M0 then M0A0)</p> <p><math>x = \sqrt[3]{\frac{120}{4\pi}}</math> or answers which round to 2.12 (-2.12 is A0)</p> <p><math>A = 2\pi(2.12)^2 + \frac{120}{2.12}, = 85</math> (only ft <math>x = 2</math> or <math>2.1</math> – both give 85)</p> <p><b>Either</b> <math>\frac{d^2 A}{dx^2} = 4\pi + \frac{240}{x^3}</math> and sign considered ( May appear in (c) )</p> <p><b>Or (method 2)</b> considers gradient to left and right of their 2.12 (e.g at 2 and 2.5)</p> <p><b>Or (method 3)</b> considers value of A either side</p> <hr/> <p>Finds numerical values for gradients and observes which is <math>&gt; 0</math> and therefore minimum (most substitute 2.12 but it is not essential to see a substitution ) (may appear in (c) )</p> <p>gradients go from negative to zero to positive so concludes minimum</p> <p><b>OR</b> finds numerical values of A , observing greater than minimum value and draws conclusion</p>	<p>B1 (1)</p> <p>B1</p> <p>M1</p> <p>A1 cso (3)</p> <p>M1 A1</p> <p>M1</p> <p>dM1 A1 (5)</p> <p>M1, A1 (2)</p> <p>M1</p> <p>A1 (2)</p> <p><b>13 marks</b></p>
<p>Notes</p>	<p>(a) <b>B1</b>: This expression must be correct and in part (a) <math>\frac{60}{\pi r^2}</math> is B0</p> <p>(b) <b>B1</b>: Accept any equivalent correct form – may be on two or more lines.</p> <p><b>M1</b> : substitute their expression for <math>h</math> in terms of <math>x</math> into Area formula <b>of the form</b> <math>kx^2 + cxh</math></p> <p><b>A1</b>: There should have been no errors in part (b) in obtaining this printed answer</p> <p>(c) <b>M1</b>: At least one power of <math>x</math> decreased by 1 <b>A1</b> accept any equivalent correct answer</p> <p><b>M1</b>: Setting <math>\frac{dA}{dx} = 0</math> and finding a value for <math>x^3</math> (<math>x^3 =</math> may be implied by answer). Allow <math>\frac{dy}{dx} = 0</math></p> <p><b>dM1</b>: Using <b>cube root</b> to find <math>x</math></p> <p><b>A1</b> : For any equivalent correct answer (need 3sf or more) Correct answer implies previous M mark</p> <p>(d) <b>M1</b> : Substitute the (+ve) <math>x</math> value found in (c) into equation for A and evaluate . <b>A1</b> is for <b>85 only</b></p> <p>(e) <b>M1</b>: Complete method, usually one of the three listed in the scheme. For first method <math>A''(x)</math> must be attempted and sign considered</p> <p><b>A1</b>: Clear statements and conclusion. (numerical substitution of <math>x</math> is not necessary in first method shown, and <math>x</math> or calculation could be wrong but <math>A''(x)</math> <b>must be correct</b> . Must <b>not</b> see 85 substituted)</p>	

Question	Scheme	Marks
<p>9 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p><math>(S_n =) a + ar + (ar^2) + \dots + ar^{n-1}</math> <b>and</b> <math>rS_n = ar + ar^2 + (ar^3) \dots + ar^n</math></p> <p><math>S_n - rS_n = a - ar^n</math></p> <p><math>S_n(1-r) = a(1-r^n)</math></p> <p>And so result <math>S_n = \frac{a(1-r^n)}{(1-r)}</math> *</p> <p>Divides one term by other (either way) to give <math>r^2 = \dots</math> then square roots to give <math>r =</math></p> <p><math>r^2 = \frac{1.944}{5.4}</math>, <math>r = 0.6</math> (ignore <math>-0.6</math>)</p> <p>Uses <math>5.4 \div r^2</math> or <math>1.944 \div r^4</math>, to give <math>a =</math></p> <p><math>a = 15</math></p> <p>Uses <math>S = \frac{15}{1-0.6}</math>, to obtain 37.5</p>	<p>M1</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>(4)</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>M1, A1ft</p> <p>(2)</p> <p>M1A1 ,A1</p> <p>(3)</p> <p><b>11 marks</b></p>
<p>Notes</p> <p>Special Case</p>	<p>(a) <b>M1:</b> Lists <b>both</b> of these sums (<math>S_n =</math>) may be omitted, <math>rS_n</math> (or <math>rS</math>) must be stated</p> <p>1<sup>st</sup> two terms must be correct in each series. Last term must be <math>ar^{n-1}</math> or <math>ar^n</math> in first series and the corresponding <math>ar^n</math> or <math>ar^{n+1}</math> in second series. Must be <math>n</math> and not a number. Reference made to other terms e.g. space or dots to indicate missing terms</p> <p><b>M1:</b> Subtracts series for <math>rS</math> from series for <math>S</math> (or other way round) to give <math>RHS = \pm(a - ar^n)</math>. This may have been obtained by following a pattern. <b>If wrong power stated on line 1 M0 here.</b> (Ignore LHS)M0M0M0A0</p> <p>d<b>M1:</b> Factorises both sides correctly– must follow from a previous M1 (It is possible to obtain M0M1M1A0 or M1M0M1A0) <b>A1:</b> completes the proof with <b>no errors</b> seen</p> <p>No errors seen: <b>First line absolutely correct</b>, omission of second line, third and fourth lines correct: M1M0M1A1</p> <p>See next sheet of common errors.</p> <p>Refer any attempts involving <b>sigma notation</b>, or any <b>proofs by induction</b> to team leader.</p> <p>Also attempts which <b>begin with the answer and work backwards</b>.</p> <p>(b) <b>M1:</b> Deduces <math>r^2</math> by dividing either term by other and attempts square root</p> <p><b>A1:</b> any correct equivalent for <math>r</math> e.g. <math>3/5</math> Answer only is <math>2/2</math></p> <p>(Method 2) Those who find fourth term must use <math>\sqrt{ab}</math> and <b>not</b> <math>\frac{1}{2}(a+b)</math> then must use it in a division with given term to obtain <math>r =</math></p> <p>(c) <b>M1:</b> May be done in two steps or more e.g. <math>5.4 \div r</math> then divided by <math>r</math> again</p> <p><b>A1ft:</b> follow through their value of <math>r</math>. Just <math>a = 15</math> <b>with no wrong working</b> implies M1A1</p> <p>(d) <b>M1:</b> States sum to infinity formula with values of <math>a</math> and <math>r</math> found earlier, provided <math> r  &lt; 1</math></p> <p><b>A1 :</b> uses 15 and 0.6 (or <math>3/5</math>) (This is <b>not</b> a ft mark) <b>A1:</b> 37.5 or exact equivalent</p>	
<p>Common errors</p>	<p>(i) Fraction inverted in (b) <math>r^2 = \frac{5.4}{1.944}</math> and <math>r = 1\frac{2}{3}</math>, then correct ft gives M1A0 M1 A1ft M0A0A0 i.e. <math>3/7</math></p> <p>(ii) Uses <math>r = 0.36</math>: (b)M0A0 (c)M1A1ft (d) M1A0A0 i.e. <math>3/7</math></p> <p>(iii) Uses <math>ar^3 = 5.4</math>, <math>ar^5 = 1.944</math> Likely to have (b)M1A1 (c)M0A0 (d) M1A0A0 i.e.<math>3/7</math></p>	

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UA031953 Summer 2012

For more information on Edexcel qualifications, please visit our website  
[www.edexcel.com](http://www.edexcel.com)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual  




Llywodraeth Cynulliad Cymru  
Welsh Assembly Government

