

Mark Scheme (Results) Summer 2008





GCE

GCE Mathematics (6665/01)

June 2008
6665 Core Mathematics C3
Mark Scheme

| Question Number | Scheme | Marks |
|-----------------|--|---|
| 1. | <p>(a)</p> $e^{2x+1} = 2$ $2x+1 = \ln 2$ $x = \frac{1}{2}(\ln 2 - 1)$ <p>(b)</p> $\frac{dy}{dx} = 8e^{2x+1}$ $x = \frac{1}{2}(\ln 2 - 1) \Rightarrow \frac{dy}{dx} = 16$ $y - 8 = 16 \left(x - \frac{1}{2}(\ln 2 - 1) \right)$ $y = 16x + 16 - 8 \ln 2$ | <p>M1</p> <p>A1 (2)</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1 (4)</p> <p>[6]</p> |

| Question Number | Scheme | Marks |
|-----------------|---|---|
| 2. | (a) $R^2 = 5^2 + 12^2$ $R = 13$ $\tan \alpha = \frac{12}{5}$ $\alpha \approx 1.176$ | M1 A1 M1 A1 (4) |
| | (b) $\cos(x - \alpha) = \frac{6}{13}$ $x - \alpha = \arccos \frac{6}{13} = 1.091 \dots$ $x = 1.091 \dots + 1.176 \dots \approx 2.267 \dots$ $x - \alpha = -1.091 \dots$ $x = -1.091 \dots + 1.176 \dots \approx 0.0849 \dots$ | M1 A1 A1 awrt 2.3 M1 accept ... = 5.19 ... for M A1 awrt 0.084 or 0.085 (5) |
| | (c)(i) $R_{\max} = 13$ (ii) At the maximum, $\cos(x - \alpha) = 1$ or $x - \alpha = 0$ $x = \alpha = 1.176 \dots$ | fit their R B1 ft M1 A1ft awrt 1.2, fit their α (3) [12] |

| Question Number | Scheme | Marks |
|-----------------|---|---|
| 3. | <p>(a)</p>  <p style="text-align: right;">  shape Vertices correctly placed </p> <p>(b)</p>  <p style="text-align: right;">  shape Vertex and intersections with axes correctly placed </p> <p>(c)</p> <p style="text-align: center;"> $P: (-1, 2)$ $Q: (0, 1)$ $R: (1, 0)$ </p> <p>(d)</p> <p> $x > -1; \quad 2 - x - 1 = \frac{1}{2}x$ Leading to $x = \frac{2}{3}$ $x < -1; \quad 2 + x + 1 = \frac{1}{2}x$ Leading to $x = -6$ </p> | <p>B1 B1 (2)</p> <p>B1 B1 (2)</p> <p>B1 B1 B1 (3)</p> <p>M1 A1 A1 M1 A1 (5) [12]</p> |

| Question Number | Scheme | Marks |
|-----------------|--|---|
| 4. | <p>(a) $x^2 - 2x - 3 = (x-3)(x+1)$</p> $f(x) = \frac{2(x-1) - (x+1)}{(x-3)(x+1)} \left(\text{or } \frac{2(x-1)}{(x-3)(x+1)} - \frac{x+1}{(x-3)(x+1)} \right)$ $= \frac{x-3}{(x-3)(x+1)} = \frac{1}{x+1} *$ | <p>B1</p> <p>M1 A1</p> <p>cso A1 (4)</p> |
| | <p>(b) $\left(0, \frac{1}{4}\right)$ Accept $0 < y < \frac{1}{4}$, $0 < f(x) < \frac{1}{4}$ etc.</p> | <p>B1 B1 (2)</p> |
| | <p>(c) Let $y = f(x)$</p> $y = \frac{1}{x+1}$ $x = \frac{1}{y+1}$ $yx + x = 1$ $y = \frac{1-x}{x}$ $f^{-1}(x) = \frac{1-x}{x}$ | <p>or $\frac{1}{x} - 1$</p> <p>M1 A1</p> |
| | <p>(d) $fg(x) = \frac{1}{2x^2 - 3 + 1}$</p> $\frac{1}{2x^2 - 2} = \frac{1}{8}$ $x^2 = 5$ $x = \pm\sqrt{5}$ | <p>ft their part (b)</p> <p>B1 ft (3)</p> <p>M1</p> <p>A1</p> <p>both A1 (3)</p> <p>[12]</p> |

| Question Number | Scheme | Marks |
|-----------------|---|--|
| 5. | <p>(a) $\sin^2 \theta + \cos^2 \theta = 1$ $\div \sin^2 \theta$ $\frac{\sin^2 \theta}{\sin^2 \theta} + \frac{\cos^2 \theta}{\sin^2 \theta} = \frac{1}{\sin^2 \theta}$ $1 + \cot^2 \theta = \operatorname{cosec}^2 \theta$ *</p> <p><i>Alternative for (a)</i> $1 + \cot^2 \theta = 1 + \frac{\cos^2 \theta}{\sin^2 \theta} = \frac{\sin^2 \theta + \cos^2 \theta}{\sin^2 \theta} = \frac{1}{\sin^2 \theta}$ $= \operatorname{cosec}^2 \theta$ *</p> <p>(b) $2(\operatorname{cosec}^2 \theta - 1) - 9 \operatorname{cosec} \theta = 3$ $2 \operatorname{cosec}^2 \theta - 9 \operatorname{cosec} \theta - 5 = 0$ or $5 \sin^2 \theta + 9 \sin \theta - 2 = 0$ $(2 \operatorname{cosec} \theta + 1)(\operatorname{cosec} \theta - 5) = 0$ or $(5 \sin \theta - 1)(\sin \theta + 2) = 0$ $\operatorname{cosec} \theta = 5$ or $\sin \theta = \frac{1}{5}$ $\theta = 11.5^\circ, 168.5^\circ$</p> | <p>M1 A1 (2) cso</p> <p>M1 A1 cso</p> <p>M1 M1 M1 A1 A1 A1 (6) [8]</p> |

| Question Number | Scheme | Marks |
|-----------------|--|--|
| 6. | (a)(i) $\frac{d}{dx}(e^{3x}(\sin x + 2 \cos x)) = 3e^{3x}(\sin x + 2 \cos x) + e^{3x}(\cos x - 2 \sin x)$ $(= e^{3x}(\sin x + 7 \cos x))$ | M1 A1 A1 (3) |
| | (ii) $\frac{d}{dx}(x^3 \ln(5x+2)) = 3x^2 \ln(5x+2) + \frac{5x^3}{5x+2}$ | M1 A1 A1 (3) |
| | (b) $\frac{dy}{dx} = \frac{(x+1)^2(6x+6) - 2(x+1)(3x^2+6x-7)}{(x+1)^4}$ $= \frac{(x+1)(6x^2+12x+6-6x^2-12x+14)}{(x+1)^4}$ $= \frac{20}{(x+1)^3} *$ | M1 $\frac{A1}{A1}$ M1 cso A1 (5) |
| | (c) $\frac{d^2y}{dx^2} = -\frac{60}{(x+1)^4} = -\frac{15}{4}$ $(x+1)^4 = 16$ $x = 1, -3$ | M1 M1 both A1 (3) |
| | <p><i>Note:</i> The simplification in part (b) can be carried out as follows</p> $\frac{(x+1)^2(6x+6) - 2(x+1)(3x^2+6x-7)}{(x+1)^4}$ $= \frac{(6x^3+18x^2+18x+6) - (6x^3+18x^2-2x-14)}{(x+1)^4}$ $= \frac{20x+20}{(x+1)^4} = \frac{20(x+1)}{(x+1)^4} = \frac{20}{(x+1)^3}$ | [14] M1 A1 |

| Question Number | Scheme | Marks |
|-----------------|--|--------------------|
| 7. | (a) $f(1.4) = -0.568 \dots < 0$ $f(1.45) = 0.245 \dots > 0$ Change of sign (and continuity) $\Rightarrow \alpha \in (1.4, 1.45)$ | M1 A1 (2) |
| | (b) $3x^3 = 2x + 6$ $x^3 = \frac{2x}{3} + 2$ $x^2 = \frac{2}{3} + \frac{2}{x}$ $x = \sqrt{\left(\frac{2}{x} + \frac{2}{3}\right)} *$ | M1 A1 A1 (3) |
| | (c) $x_1 = 1.4371$ $x_2 = 1.4347$ $x_3 = 1.4355$ | B1 B1 B1 (3) |
| | (d) Choosing the interval (1.4345, 1.4355) or appropriate tighter interval. $f(1.4345) = -0.01 \dots$ $f(1.4355) = 0.003 \dots$ Change of sign (and continuity) $\Rightarrow \alpha \in (1.4345, 1.4355)$ | M1 M1 |
| | $\Rightarrow \alpha = 1.435$, correct to 3 decimal places * cso Note: $\alpha = 1.435\ 304\ 553 \dots$ | A1 (3) [11] |