

Question 1 continued

Blank lined area for writing the answer to Question 1 continued.

Leave blank

Q1

(Total 5 marks)



Leave
blank

3.




Figure 1

A small box is pushed along a floor. The floor is modelled as a rough horizontal plane and the box is modelled as a particle. The coefficient of friction between the box and the floor is $\frac{1}{2}$. The box is pushed by a force of magnitude 100 N which acts at an angle of 30° with the floor, as shown in Figure 1.

Given that the box moves with constant speed, find the mass of the box.

(7)

Handwritten answer area with horizontal lines.





Question 4 continued

Leave blank



Handwriting practice area consisting of 30 horizontal lines.



N 3 5 3 9 0 A 0 9 2 8



Question 6 continued

Leave blank

Lined area for writing answers.



Question 6 continued

Lined area for writing the answer to Question 6.

Leave blank

Q6

(Total 10 marks)



Leave blank

7.




Figure 2

A particle of mass 0.4 kg is held at rest on a fixed rough plane by a horizontal force of magnitude P newtons. The force acts in the vertical plane containing the line of greatest slope of the inclined plane which passes through the particle. The plane is inclined to the horizontal at an angle α , where $\tan \alpha = \frac{3}{4}$, as shown in Figure 2.

The coefficient of friction between the particle and the plane is $\frac{1}{3}$.

Given that the particle is on the point of sliding up the plane, find

(a) the magnitude of the normal reaction between the particle and the plane, (5)

(b) the value of P . (5)





Question 7 continued

Leave blank

Lined area for writing answers.



Question 7 continued

(This area contains 30 horizontal lines for the student to write their answer to Question 7.)

(Total 10 marks)

Leave blank

Q7



Leave blank

8.




Figure 3

Two particles A and B have mass 0.4 kg and 0.3 kg respectively. The particles are attached to the ends of a light inextensible string. The string passes over a small smooth pulley which is fixed above a horizontal floor. Both particles are held, with the string taut, at a height of 1 m above the floor, as shown in Figure 3. The particles are released from rest and in the subsequent motion B does not reach the pulley.

(a) Find the tension in the string immediately after the particles are released. (6)

(b) Find the acceleration of A immediately after the particles are released. (2)

When the particles have been moving for 0.5 s , the string breaks.

(c) Find the further time that elapses until B hits the floor. (9)



