


1.

Figure 1



A particle P is attached to one end of a light inextensible string. The other end of the string is attached to a fixed point O . A horizontal force of magnitude 12 N is applied to P . The particle P is in equilibrium with the string taut and OP making an angle of 20° with the downward vertical, as shown in Figure 1.

Find

- (a) the tension in the string, (3)

- (b) the weight of P . (4)



Leave
blank

Question 1 continued

Lined writing area for Question 1 continued.

(Total 7 marks)

Q1



Question 3 continued

Leave
blank

Lined area for writing the answer to Question 3.

(Total 9 marks)

Q3



Leave
blank

4. A car is moving along a straight horizontal road. At time $t=0$, the car passes a point A with speed 25 m s^{-1} . The car moves with constant speed 25 m s^{-1} until $t=10 \text{ s}$. The car then decelerates uniformly for 8 s . At time $t=18 \text{ s}$, the speed of the car is $V \text{ m s}^{-1}$ and this speed is maintained until the car reaches the point B at time $t=30 \text{ s}$.

(a) Sketch, in the space below, a speed–time graph to show the motion of the car from A to B . (3)

Given that $AB = 526 \text{ m}$, find

(b) the value of V , (5)

(c) the deceleration of the car between $t=10 \text{ s}$ and $t=18 \text{ s}$. (3)



Question 4 continued

Lined writing area for the answer to Question 4.

Leave
blank




N 2 6 1 1 4 A 0 9 2 0

Leave blank

6.

Figure 4



Two particles P and Q have mass 0.5 kg and $m \text{ kg}$ respectively, where $m < 0.5$. The particles are connected by a light inextensible string which passes over a smooth, fixed pulley. Initially P is 3.15 m above horizontal ground. The particles are released from rest with the string taut and the hanging parts of the string vertical, as shown in Figure 4. After P has been descending for 1.5 s , it strikes the ground. Particle P reaches the ground before Q has reached the pulley.

- (a) Show that the acceleration of P as it descends is 2.8 m s^{-2} . (3)
- (b) Find the tension in the string as P descends. (3)
- (c) Show that $m = \frac{5}{18}$. (4)
- (d) State how you have used the information that the string is inextensible. (1)

When P strikes the ground, P does not rebound and the string becomes slack. Particle Q then moves freely under gravity, without reaching the pulley, until the string becomes taut again.

- (e) Find the time between the instant when P strikes the ground and the instant when the string becomes taut again. (6)



Question 6 continued

Lined area for writing answers to Question 6.

Leave blank

Q6

(Total 17 marks)



Leave
blank

7. A boat B is moving with constant velocity. At noon, B is at the point with position vector $(3\mathbf{i} - 4\mathbf{j})$ km with respect to a fixed origin O . At 1430 on the same day, B is at the point with position vector $(8\mathbf{i} + 11\mathbf{j})$ km.

(a) Find the velocity of B , giving your answer in the form $p\mathbf{i} + q\mathbf{j}$. (3)

At time t hours after noon, the position vector of B is \mathbf{b} km.

(b) Find, in terms of t , an expression for \mathbf{b} . (3)

Another boat C is also moving with constant velocity. The position vector of C , \mathbf{c} km, at time t hours after noon, is given by

$$\mathbf{c} = (-9\mathbf{i} + 20\mathbf{j}) + t(6\mathbf{i} + \lambda\mathbf{j}),$$

where λ is a constant. Given that C intercepts B ,

(c) find the value of λ , (5)

(d) show that, before C intercepts B , the boats are moving with the same speed. (3)



Leave blank

Question 7 continued

Lined writing area for the answer to Question 7 continued.



N 2 6 1 1 4 A 0 1 9 2 0

