



Mark Scheme (Results)

Summer 2014

Pearson Edexcel GCE in Core Mathematics 2  
(6664\_01)

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2014

Publications Code UA038455

All the material in this publication is copyright

© Pearson Education Ltd 2014

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## PEARSON EDEXCEL GCE MATHEMATICS

### General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - d... or dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper or ag- answer given
  - $\square$  or d... The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
  - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

## General Principles for Core Mathematics Marking

*(But note that specific mark schemes may sometimes override these general principles).*

### **Method mark for solving 3 term quadratic:**

#### **1. Factorisation**

$(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$ , leading to  $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$ , leading to  $x = \dots$

#### **2. Formula**

Attempt to use the correct formula (with values for a, b and c).

#### **3. Completing the square**

Solving  $x^2 + bx + c = 0$ :  $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$ ,  $q \neq 0$ , leading to  $x = \dots$

### **Method marks for differentiation and integration:**

#### **1. Differentiation**

Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )

#### **2. Integration**

Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

### **Use of a formula**

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

### **Exact answers**

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

| Question Number   | Scheme  |  |   |        |              | Marks                 |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|---|---|--|---|--------|--------------|-----------------------|-------|---|-------|------|-------|-----|--|------|--|---|-----|--|-------|--|--------------|--|-------|--|-------|--|-------|--|
| 1.(a)   | <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="border: none;"><math>x</math></td> <td style="border: none;">  </td> <td style="border: none;">1</td> <td style="border: none;"> </td> <td style="border: none;">1.25</td> <td style="border: none;"> </td> <td style="border: none;">1.5</td> <td style="border: none;"> </td> <td style="border: none;">1.75</td> <td style="border: none;"> </td> <td style="border: none;">2</td> </tr> <tr> <td style="border: none;"><math>y</math></td> <td style="border: none;">  </td> <td style="border: none;">1.414</td> <td style="border: none;"> </td> <td style="border: none;"><b>1.601</b></td> <td style="border: none;"> </td> <td style="border: none;">1.803</td> <td style="border: none;"> </td> <td style="border: none;">2.016</td> <td style="border: none;"> </td> <td style="border: none;">2.236</td> </tr> </table> |  |   |        |              | $x$                   |       | 1 |       | 1.25 |       | 1.5 |  | 1.75 |  | 2 | $y$ |  | 1.414 |  | <b>1.601</b> |  | 1.803 |  | 2.016 |  | 2.236 |  |
|   | $x$   |  | 1   |        | 1.25         |                       | 1.5   |   | 1.75  |      | 2     |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|   | $y$   |  | 1.414   |        | <b>1.601</b> |                       | 1.803 |   | 2.016 |      | 2.236 |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
| {At $x = 1.25,$ } $y = 1.601$ (only)  | 1.601 ( <b>May not be in the table and can score if seen as part of their working in (b)</b> )  |  |   | B1 cao |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|   |   |  |   |        |              | <b>[1]</b>            |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
| (b)   | $\frac{1}{2} \times 0.25; \times \{1.414 + 2.236 + 2(\text{their } 1.601 + 1.803 + 2.016)\}$  |  |   |        |              | B1;<br><u>M1 A1ft</u> |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|   | B1; for using $\frac{1}{2} \times 0.25$ or $\frac{1}{8}$ or equivalent.   | <u>M1: Structure of</u><br>$\{.....\}$ | <u>A1ft:</u> for the correct expression as shown following through candidate's $y$ value found in part (a). |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|   | <p>M1 requires the correct structure for the <math>y</math> values. It needs to contain first <math>y</math> value <b>plus</b> last <math>y</math> value and the second bracket to be multiplied by 2 and to be the summation of the remaining <math>y</math> values in the table with no additional values. If the only mistake is a copying error or is to omit one value from 2(.....) bracket this may be regarded as a slip and the M mark can be allowed (nb: an extra repeated term, however, forfeits the M mark). M0 if any values used are <math>x</math> values instead of <math>y</math> values.</p> <p>A1ft: for the correct underlined expression as shown following through candidate's <math>y</math> value found in part (a).</p> <p>Bracketing mistakes: e.g.</p>   |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|   | $\left(\frac{1}{2} \times \frac{1}{4}\right)(1.414 + 2.236) + 2(\text{their } 1.601 + 1.803 + 2.016)(=11.29625)$  |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|   | $\left(\frac{1}{2} \times \frac{1}{4}\right)1.414 + 2.236 + 2(\text{their } 1.601 + 1.803 + 2.016)(=13.25275)$  |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|   | <p>Both score B1 M1 A0 unless the final answer implies that the calculation has been done correctly (then full marks could be given).</p>   |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
|   | <p><b>Alternative:</b><br/>Separate trapezia may be used, and this can be marked equivalently.</p> $\left[ \frac{1}{8}(1.414 + 1.601) + \frac{1}{8}(1.601 + 1.803) + \frac{1}{8}(1.803 + 2.016) + \frac{1}{8}(2.016 + 2.236) \right]$ <p>B1 for <math>\frac{1}{8}</math> (aef), M1 for correct structure, 1st A1ft for correct expression, ft their 1.601</p>   |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
| $\left\{ = \frac{1}{8}(14.49) \right\} = 1.81125$   |   |  | 1.81 or awrt 1.81   |        | A1           |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
| <b>Correct answer <u>only</u> in (b) scores no marks</b>  |   |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
| <b>If required accuracy is not seen in (a), full marks can still be scored in (b) (e.g. uses 1.6)</b> |   |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
| <b>[4]</b>  |   |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |
| <b>Total 5</b>  |   |  |   |        |              |                       |       |   |       |      |       |     |  |      |  |   |     |  |       |  |              |  |       |  |       |  |       |  |



| Question Number | Scheme   |   | Marks          |
|-----------------|--|---|----------------|
|                 | <b>If there is no labelling, mark (a) and (b) in that order</b>  |   |                |
|                 | $f(x) = 2x^3 - 7x^2 + 4x + 4$  |   |                |
| 2.(a)           | $f(2) = 2(2)^3 - 7(2)^2 + 4(2) + 4$  | Attempts $f(2)$ or $f(-2)$  | M1             |
|                 | $= 0$ , and so $(x - 2)$ is a factor.  | $f(2) = 0$ with no sign or substitution errors<br>$(2(2)^3 - 7(2)^2 + 4(2) + 4 = 0$ is sufficient)<br><b>and for conclusion.</b> Stating “hence factor” or “it is a factor” or a “tick” or “QED” or “no remainder” or “as required” are fine for the conclusion <b>but not = 0 just underlined and not hence (2 or f(2)) is a factor.</b> Note also that a conclusion can be implied from a <u>preamble</u> , eg: “If $f(2) = 0$ , $(x - 2)$ is a factor....” | A1             |
|                 | <b>Note: Long division scores no marks in part (a). The <u>factor theorem</u> is required.</b>   |   |                |
|                 |  |   | <b>[2]</b>     |
| (b)             | $f(x) = \{(x - 2)\}(2x^2 - 3x - 2)$  | M1: Attempts long division by $(x - 2)$ or other method using $(x - 2)$ , to obtain $(2x^2 \pm ax \pm b)$ , $a \neq 0$ , even with a remainder. Working need not be seen as this could be done “by inspection.”<br>A1: $(2x^2 - 3x - 2)$  | M1 A1          |
|                 | $= (x - 2)(x - 2)(2x + 1)$ or $(x - 2)^2(2x + 1)$<br>or equivalent e.g.<br>$= 2(x - 2)(x - 2)(x + \frac{1}{2})$ or $2(x - 2)^2(x + \frac{1}{2})$ | <b>dM1:</b> Factorises a 3 term quadratic. (see rule for factorising a quadratic in the General Principles for Core Maths Marking). <b>This is dependent</b> on the previous method mark being awarded but there must have been no remainder. Allow an attempt to solve the quadratic to determine the <b>factors</b> .<br>A1: cao – needs all three factors <b>on one line</b> . Ignore following work (such as a solution to a quadratic equation.)         | dM1 A1         |
|                 | Note $= (x - 2)(\frac{1}{2}x - 1)(4x + 2)$ would lose the last mark as it is not <b>fully</b> factorised   |   |                |
|                 | <b>For correct answers only award full marks in (b)</b>  |   |                |
|                 |  |   | <b>[4]</b>     |
|                 |  |   | <b>Total 6</b> |


| Question Number | Scheme   |   | Marks          |
|-----------------|--|---|----------------|
| 3. (a)          | $(2 - 3x)^6 = 64 + \dots$  | 64 seen as the only constant term in their expansion.   | B1             |
|                 | $\{(2 - 3x)^6\} = (2)^6 + {}^6C_1(2)^5(-3x) + {}^6C_2(2)^4(-3x)^2 + \dots$   |   | M1             |
|                 | M1: $({}^6C_1 \times \dots \times x)$ or $({}^6C_2 \times \dots \times x^2)$ . For <u>either</u> the $x$ term <u>or</u> the $x^2$ term. Requires <u>correct</u> binomial coefficient in any form <u>with the correct power of <math>x</math></u> , but the other part of the coefficient (perhaps including powers of 2 and/or $-3$ ) may be wrong or missing. The terms can be “listed” rather than added. Ignore any extra terms.  |   |                |
|                 | ${}^6C_1 2^5 - 3x + {}^6C_2 2^4 - 3x^2 + \dots$ Scores M0 unless later work implies a correct method   |   |                |
|                 | $= 64 - 576x + 2160x^2 + \dots$  | A1: Either $-576x$ or $2160x^2$<br>(Allow $+ -576x$ here)<br>A1: Both $-576x$ and $2160x^2$<br>(Do not allow $+ -576x$ here)  | A1A1           |
|                 |  | [4]   |                |
| (a) Way 2       | $(2 - 3x)^6 = 64 + \dots$  | 64 seen as the only constant term in their expansion.   | B1             |
|                 | $\left(1 - \frac{3}{2}x\right)^6 = 1 + {}^6C_1\left(\frac{-3}{2}x\right) + {}^6C_2\left(\frac{-3}{2}x\right)^2 + \dots$  | M1: $({}^6C_1 \times \dots \times x)$ or $({}^6C_2 \times \dots \times x^2)$ . For <u>either</u> the $x$ term <u>or</u> the $x^2$ term. Requires <u>correct</u> binomial coefficient in any form <u>with the correct power of <math>x</math></u> , but the other part of the coefficient (perhaps including powers of 2 and/or $-3$ ) may be wrong or missing. The terms can be “listed” rather than added. Ignore any extra terms. | M1             |
|                 | $= 64 - 576x + 2160x^2 + \dots$  | A1: Either $-576x$ or $2160x^2$<br>(Allow $+ -576x$ here)<br>A1: Both $-576x$ and $2160x^2$<br>(Do not allow $+ -576x$ here)  | A1A1           |
| (b)             | Candidate writes down $\left(1 + \frac{x}{2}\right) \times$ (their part (a) answer, at least up to the term in $x$ ).<br>(Condone missing brackets)<br>$\left(1 + \frac{x}{2}\right)(64 - 576x + \dots)$ or $\left(1 + \frac{x}{2}\right)(64 - 576x + 2160x^2 + \dots)$ or<br>$\left(1 + \frac{x}{2}\right)64 - \left(1 + \frac{x}{2}\right)576x$ or $\left(1 + \frac{x}{2}\right)64 - \left(1 + \frac{x}{2}\right)576x + \left(1 + \frac{x}{2}\right)2160x^2$<br>or $64 + 32x, -576x - 288x^2, 2160x^2 + 1080x^3$ are fine. |   | M1             |
|                 | $= 64 - 544x + 1872x^2 + \dots$  | A1: At least 2 terms correct as shown.<br>(Allow $+ -544x$ here)<br>A1: $64 - 544x + 1872x^2$<br>The terms can be “listed” rather than added. Ignore any extra terms.   | A1A1           |
|                 |  |   | [3]            |
|                 |  |   | <b>Total 7</b> |
|                 | <b>SC: If a candidate expands in descending powers of <math>x</math>, only the M marks are available</b>   |   |                |
|                 | e.g. $\{(2 - 3x)^6\} = (-3x)^6 + {}^6C_1(2)^5(-3x)^5 + {}^6C_2(2)^4(-3x)^4 + \dots$  |   |                |

| Question Number   | Scheme   |   | Marks                        |            |
|---|--|---|------------------------------|------------|
| <p><b>4.</b></p> $\left\{ \int \left( \frac{x^3}{6} + \frac{1}{3x^2} \right) dx \right\} = \frac{x^4}{6(4)} + \frac{x^{-1}}{(3)(-1)}$   |  | <p>M1: <math>x^n \rightarrow x^{n+1}</math></p>   | <p>M1A1A1</p>                |            |
|   |  | <p>A1: At least one of either <math>\frac{x^4}{6(4)}</math> or <math>\frac{x^{-1}}{(3)(-1)}</math>.</p>   |                              |            |
|   |  | <p>A1: <math>\frac{x^4}{6(4)} + \frac{x^{-1}}{(3)(-1)}</math> or equivalent.<br/>                     e.g. <math>\frac{x^4}{6} + \frac{x^{-1}}{-3}</math> (they will lose the final mark if they cannot deal with this correctly)</p> |                              |            |
|   | <p><b>Note that some candidates may change the function prior to integrating e.g.</b><br/> <math>\int \frac{x^3}{6} + \frac{1}{3x^2} dx = \int 3x^5 + 6 dx</math> in which case allow the M1 if <math>x^n \rightarrow x^{n+1}</math> for their changed function and allow the M1 for limits if scored</p>  |   |                              |            |
|   | $\left\{ \int_1^{\sqrt{3}} \left( \frac{x^3}{6} + \frac{1}{3x^2} \right) dx \right\} = \left( \frac{(\sqrt{3})^4}{24} + \frac{(\sqrt{3})^{-1}}{-1(3)} \right) - \left( \frac{(1)^4}{24} + \frac{(1)^{-1}}{-1(3)} \right)$  |   |                              | <p>dM1</p> |
| <p>2<sup>nd</sup> dM1: For using limits of <math>\sqrt{3}</math> and 1 on an integrated expression and subtracting the correct way round. <b>The 2<sup>nd</sup> M1 is dependent on the 1<sup>st</sup> M1 being awarded.</b></p> |  |   |                              |            |
| $= \left( \frac{9}{24} - \frac{1}{3\sqrt{3}} \right) - \left( \frac{1}{24} - \frac{1}{3} \right) = \frac{2}{3} - \frac{1}{9}\sqrt{3}$   | $\frac{2}{3} - \frac{1}{9}\sqrt{3}$ or $a = \frac{2}{3}$ and $b = -\frac{1}{9}$ .<br>Allow equivalent fractions for $a$ and/or $b$ and 0.6 recurring and/or 0.1 recurring but do <b>not</b> allow $\frac{6-\sqrt{3}}{9}$   | <p>A1cso</p>  |                              |            |
| <p><b>This final mark is cao and cso – there must have been no previous errors</b></p>  |  |   |                              |            |
|   |  |   | <p><b>Total 5</b></p>        |            |
| <p><b>Common Errors (Usually 3 out of 5)</b></p>  |  |   |                              |            |
|   | $\left\{ \int \left( \frac{x^3}{6} + \frac{1}{3x^2} \right) dx \right\} = \int \left( \frac{x^3}{6} + 3x^{-2} \right) dx = \frac{x^4}{6(4)} + \frac{3x^{-1}}{(-1)}$ $\left\{ \int_1^{\sqrt{3}} \left( \frac{x^3}{6} + \frac{1}{3x^2} \right) dx \right\} = \left( \frac{(\sqrt{3})^4}{24} + \frac{3(\sqrt{3})^{-1}}{-1} \right) - \left( \frac{(1)^4}{24} + \frac{3(1)^{-1}}{-1} \right)$ $= \left( \frac{9}{24} - \frac{3}{\sqrt{3}} \right) - \left( \frac{1}{24} + \frac{3}{-1} \right) = \frac{10}{3} - \sqrt{3}$                      |   | <p>M1A1A0<br/>dM1<br/>A0</p> |            |
|   | $\left\{ \int \left( \frac{x^3}{6} + \frac{1}{3x^2} \right) dx \right\} = \int \left( \frac{x^3}{6} + (3x)^{-2} \right) dx = \frac{x^4}{6(4)} + \frac{(3x)^{-1}}{(-1)}$ $\left\{ \int_1^{\sqrt{3}} \left( \frac{x^3}{6} + \frac{1}{3x^2} \right) dx \right\} = \left( \frac{(\sqrt{3})^4}{24} + \frac{(3\sqrt{3})^{-1}}{-1} \right) - \left( \frac{(1)^4}{24} + \frac{(3 \times 1)^{-1}}{-1} \right)$ $= \left( \frac{9}{24} - \frac{1}{3\sqrt{3}} \right) - \left( \frac{1}{24} - \frac{1}{3} \right) = \frac{2}{3} - \frac{\sqrt{3}}{9}$ |   | <p>M1A1A0<br/>dM1<br/>A0</p> |            |
| <p><b>Note this is the correct answer but follows incorrect work.</b></p>   |  |   |                              |            |

| Question Number  | Scheme   |  | Marks |
|--|--|--|-------|
| 5.(a)  | $\text{Area } BDE = \frac{1}{2}(5)^2(1.4)$   | M1: Use of the correct formula or method for the area of the sector  | M1A1  |
|  | $= 17.5 \text{ (cm}^2\text{)}$   | A1: 17.5 oe  |       |
|  |  |  | [2]   |
| (b)  | <b>Parts (b) and (c) can be marked together</b>  |  |       |
|  | $6.1^2 = 5^2 + 7.5^2 - (2 \times 5 \times 7.5 \cos DBC)$ or $\cos DBC = \frac{5^2 + 7.5^2 - 6.1^2}{2 \times 5 \times 7.5}$ (or equivalent)                         |  | M1    |
|  | M1: A correct statement involving the angle $DBC$  |  |       |
|  | Angle $DBC = 0.943201\dots$  | awrt 0.943   | A1    |
|  | <b>Note that work for (b) may be seen on the diagram or in part (c)</b>  |  |       |
|  |  | [2]  |       |
| (c)  | <b>Note that candidates may work in degrees in (c) (Angle <math>DBC = 54.04\dots</math>degrees )</b>   |  |       |
|  | $\text{Area } CBD = \frac{1}{2}5(7.5)\sin(0.943)$  |  |       |
|  | Angle $EBA = \pi - 1.4 - "0.943"$<br>(Maybe seen on the diagram)   | Area $CBD = \frac{1}{2}5(7.5)\sin(\text{their } 0.943)$ or awrt 15.2. (Note area of $CBD = 15.177\dots$ )<br>A correct method for the area of triangle $CBD$ which can be implied by awrt 15.2   | M1    |
|  | $\pi - 1.4 - \text{"their } 0.943"$  |  |       |
|  | A value for angle $EBA$ of awrt 0.8 (from 0.7985926536... or 0.7983916536...) or value for angle $EBA$ of (1.74159... – their angle $DBC$ ) would imply this mark. |  | M1    |
|  | $AB = 5 \cos(\pi - 1.4 - "0.943")$<br>or<br>$AE = 5 \sin(\pi - 1.4 - "0.943")$   |  |       |
|  |  | $AB = 5 \cos(\pi - 1.4 - \text{their } 0.943)$<br>$AB = 5 \cos(0.79859\dots) = 3.488577938\dots$<br>Allow M1 for $AB =$ awrt 3.49<br>Or<br>$AE = 5 \sin(\pi - 1.4 - \text{their } 0.943)$<br>$AE = 5 \sin(0.79859\dots) = 3.581874365688\dots$<br>Allow M1 for $AE =$ awrt 3.58<br><b>It must be clear that <math>\pi - 1.4 - "0.943"</math> is being used for angle <math>EBA</math>.</b><br><b>Note that some candidates use the sin rule here but it must be used correctly – do not allow mixing of degrees and radians.</b> | M1    |
| $\text{Area } EAB = \frac{1}{2}5 \cos(\pi - 1.4 - "0.943") \times 5 \sin(\pi - 1.4 - "0.943")$ |  |  |       |
|  | <b><u>This is dependent on the previous M1</u></b><br><b><u>and there must be no other errors in finding the area of triangle EAB</u></b>                          | dM1  |       |
|  | Allow M1 for area $EAB =$ awrt 6.2   |  |       |
|  | Area $ABCDE = 15.17\dots + 17.5 + 6.24\dots = 38.92\dots$  |  |       |
|  | awrt 38.9  | A1cso  |       |
|  |  | [5]  |       |
|  | <b>Note that a sign error in (b) can give the obtuse angle (2.198....) and could lead to the correct answer in (c) – this would lose the final mark in (c)</b>     | <b>Total</b><br><b>9</b>   |       |

| Question Number  | Scheme  |  | Marks      |
|--|---|--|------------|
| <b>6(a)</b>  | $S_{\infty} = \frac{20}{1 - \frac{7}{8}} ; = 160$   | M1: Use of a correct $S_{\infty}$ formula  | M1A1       |
|  |   | A1: 160  |            |
|  | <b>Accept correct answer only (160)</b>   |  | <b>[2]</b> |
| <b>(b)</b>   | $S_{12} = \frac{20(1 - (\frac{7}{8})^{12})}{1 - \frac{7}{8}} ; = 127.77324...$  | M1: Use of a correct $S_n$ formula with $n = 12$<br>(condone missing brackets around 7/8)  | M1A1       |
|  |   | A1: awrt 127.8   |            |
|  | <b>T &amp; I in (b) requires all 12 terms to be calculated correctly for M1 and A1 for awrt 127.8</b>   |  | <b>[2]</b> |
| <b>(c)</b>   | $160 - \frac{20(1 - (\frac{7}{8})^N)}{1 - \frac{7}{8}} < 0.5$   | Applies $S_N$ ( <b>GP only</b> ) with $a = 20$ , $r = \frac{7}{8}$ and “uses” 0.5 and their $S_{\infty}$ at any point in their working. (condone missing brackets around 7/8)(Allow =, <, >, ≥, ≤ ) but see note below.  | M1         |
|  | $160(\frac{7}{8})^N < (0.5)$ or $(\frac{7}{8})^N < (\frac{0.5}{160})$   | Attempt to isolate $+160(\frac{7}{8})^N$ or $(\frac{7}{8})^N$ oe<br>(Allow =, <, >, ≥, ≤ ) but see note below.<br><b>Dependent on the previous M1</b>  | dM1        |
|  | $N \log(\frac{7}{8}) < \log(\frac{0.5}{160})$   | Uses the power law of logarithms or takes logs base 0.875 correctly to obtain an equation or an inequality of the form<br>$N \log(\frac{7}{8}) < \log(\frac{0.5}{\text{their } S_{\infty}})$<br>or<br>$N > \log_{0.875}(\frac{0.5}{\text{their } S_{\infty}})$<br>(Allow =, <, >, ≥, ≤ ) but see note below. | M1         |
|  | $N > \frac{\log(\frac{0.5}{160})}{\log(\frac{7}{8})} = 43.19823... \Rightarrow N = 44$  | $N = 44$ (Allow $N \geq 44$ but not $N > 44$ )   | A1 cso     |
|  | An incorrect <b>inequality</b> statement at any stage in a candidate’s working loses the final mark. Some candidates do not realise that the direction of the inequality is reversed in the final line of their solution. <b>BUT</b> it is possible to gain full marks for using =, as long as no incorrect working seen. |  |            |
| <b>Total 8</b>   |   |  |            |
| <b>Trial &amp; Improvement Method in (c):</b>  |   |  |            |
| 1 <sup>st</sup> M1: Attempts $160 - S_N$ or $S_N$ with at least one value for $N > 40$   |   |  |            |
| 2 <sup>nd</sup> M1: Attempts $160 - S_N$ or $S_N$ with $N = 43$ or $N = 44$  |   |  |            |
| 3 <sup>rd</sup> M1: For evidence of examining $160 - S_N$ or $S_N$ for <b>both</b> $N = 43$ <b>and</b> $N = 44$ with <b>both</b> values correct to 2 DP<br>Eg: $160 - S_{43} = \text{awrt } 0.51$ and $160 - S_{44} = \text{awrt } 0.45$<br>or $S_{43} = \text{awrt } 159.49$ and $S_{44} = \text{awrt } 159.55$ |   |  |            |
| A1: $N = 44$ <b>cso</b>  |   |  |            |
| <b>Answer of <math>N = 44</math> only with no working scores no marks</b>  |   |  |            |

| Question Number  | Scheme  | Marks   |       |
|--|---|---|-------|
| 7.   | (i) $9\sin(\theta + 60^\circ) = 4$ ; $0 \leq \theta < 360^\circ$<br>(ii) $2\tan x - 3\sin x = 0$ ; $-\pi \leq x < \pi$  |   |       |
| (i)  | $\sin(\theta + 60^\circ) = \frac{4}{9}$ , so $(\theta + 60^\circ) = 26.3877\dots$<br><br>$(\alpha = 26.3877\dots)$  | Sight of $\sin^{-1}\left(\frac{4}{9}\right)$ or awrt $26.4^\circ$ or $0.461^\circ$<br><br>Can also be implied for $\theta =$ awrt $-33.6$ (i.e. $26.4 - 60$ )   | M1    |
|  | So, $\theta + 60^\circ = \{153.6122\dots, 386.3877\dots\}$  | <b><math>\theta + 60^\circ</math></b> = either "180 – their $\alpha$ " or "360 + their $\alpha$ " and not for $\theta =$ either "180 – their $\alpha$ " or "360 + their $\alpha$ ". This can be implied by later working. The candidate's $\alpha$ could also be in radians but do not allow mixing of degrees and radians. | M1    |
|  | and $\theta = \{93.6122\dots, 326.3877\dots\}$  | A1: At least one of<br>awrt $93.6^\circ$ or awrt $326.4^\circ$  | A1 A1 |
|  |   | A1: Both awrt $93.6^\circ$ and awrt $326.4^\circ$   |       |
|  | <b>Both answers are cso and must come from correct work</b>   |   |       |
|  | Ignore extra solutions outside the range.<br>In an otherwise fully correct solution deduct the final A1 for any extra solutions in range                        |   |       |
|  |   | <b>[4]</b>  |       |
| (ii)   | $2\left(\frac{\sin x}{\cos x}\right) - 3\sin x = 0$   | Applies $\tan x = \frac{\sin x}{\cos x}$  | M1    |
|  | Note: Applies $\tan x = \frac{\sin x}{\cos x}$ can be implied by $2\tan x - 3\sin x = 0 \Rightarrow \tan x(2 - 3\cos x)$  |   |       |
|  | $2\sin x - 3\sin x \cos x = 0$  |   |       |
|  | $\sin x(2 - 3\cos x) = 0$   |   |       |
|  | $\cos x = \frac{2}{3}$  | $\cos x = \frac{2}{3}$  | A1    |
| $x =$ awrt $\{0.84, -0.84\}$   | A1: One of either awrt $0.84$ or awrt $-0.84$<br>A1ft: You can apply ft for $x = \pm \alpha$ , where $\alpha = \cos^{-1} k$ and $-1 \leq k \leq 1$              | A1A1ft  |       |
| <b>In this part of the solution, if there are any extra answers in range in an otherwise correct solution withhold the A1ft.</b> |   |   |       |
| $\{\sin x = 0 \Rightarrow\} x = 0$ and $-\pi$  | <b>Both <math>x = 0</math> and <math>-\pi</math></b> or awrt $-3.14$ from $\sin x = 0$<br><b>In this part of the solution, ignore extra solutions in range.</b> | B1  |       |
| Note solutions are: $x = \{-3.1415\dots, -0.8410\dots, 0, 0.8410\dots\}$<br>Ignore extra solutions outside the range             |   |   |       |
| For <b>all</b> answers in degrees in (ii) M1A1A0A1ftB0 is possible   |   |   |       |
| <b>Allow the use of <math>\theta</math> in place of <math>x</math> in (ii)</b>   |   |   |       |
|  |   | <b>[5]</b>  |       |
|  |   | <b>Total 9</b>  |       |

| Question Number | Scheme  | Marks   |            |  |
|-----------------|---|---|------------|--|
| <b>8.</b>       | Graph of $y = 3^x$ and solving $3^{2x} - 9(3^x) + 18 = 0$                                       |   |            |  |
| <b>(a)</b>      |                | At least two of the three criteria correct.<br><b>(See notes below.)</b>  | B1         |  |
|                 |   | All three criteria correct.<br><b>(See notes below.)</b>  | B1         |  |
|                 |   | <b>Criteria number 1:</b> Correct shape of curve for $x \geq 0$ and at least touches the positive y-axis.<br><b>Criteria number 2:</b> Correct shape of curve for $x < 0$ . Must not touch the x-axis or have any turning points.<br><b>Criteria number 3:</b> (0, 1) stated or in a table or 1 marked on the y-axis. Allow (1, 0) rather than (0, 1) if marked in the "correct" place on the y-axis. |            |  |
|                 |   |   | <b>[2]</b> |  |
| <b>(b)</b>      | $(3^x)^2 - 9(3^x) + 18 = 0$<br>or<br>$y = 3^x \Rightarrow y^2 - 9y + 18 = 0$                    | Forms a quadratic of the correct form in $3^x$ or in "y" where "y" = $3^x$ or even in x where "x" = $3^x$   | M1         |  |
|                 | $\{ (y-6)(y-3) = 0 \text{ or } (3^x - 6)(3^x - 3) = 0 \}$                                       |   |            |  |
|                 | $y = 6, y = 3 \text{ or } 3^x = 6, 3^x = 3$   | <b>Both</b> $y = 6$ and $y = 3$ .   | A1         |  |
|                 | $\{3^x = 6 \Rightarrow\} x \log 3 = \log 6$<br>or $x = \frac{\log 6}{\log 3}$ or $x = \log_3 6$ | A valid method for solving $3^x = k$ where $k > 0, k \neq 1, k \neq 3$<br><br><b>to give either</b> <div style="border: 1px solid black; padding: 5px; display: inline-block;"> <math>x \log 3 = \log k \text{ or } x = \frac{\log k}{\log 3} \text{ or } x = \log_3 k</math> </div>  | dM1        |  |
|                 | $x = 1.63092\dots$  | awrt 1.63   | A1cso      |  |
|                 | Provided the first M1A1 is scored, the second M1A1 can be implied by awrt 1.63                  |   |            |  |
|                 | $x = 1$   | $x = 1$ stated as a solution from <b>any</b> working.   | B1         |  |
|                 | <b>[5]</b>  |   |            |  |
|                 |   | <b>Total 7</b>  |            |  |

| Question Number  | Scheme  |  | Marks      |
|--|---|--|------------|
| <b>Mark (a) and (b) together</b>   |   |  |            |
| <b>9. (a)</b>  | $OQ^2 = (6\sqrt{5})^2 + 4^2$ or $OQ = \sqrt{(6\sqrt{5})^2 + 4^2} \quad \{= 14\}$                | Uses the addition form of Pythagoras on $6\sqrt{5}$ and 4. Condone missing brackets on $(6\sqrt{5})^2$<br><b>(Working or 14 may be seen on the diagram)</b>  | M1         |
|  | $y_Q = \sqrt{14^2 - 11^2}$  | $y_Q = \sqrt{(\text{their } OQ)^2 - 11^2}$<br><b>Must include <math>\sqrt{\quad}</math> and is dependent on the first M1 and requires <math>OQ &gt; 11</math></b>  |            |
|  | $= \sqrt{75}$ or $5\sqrt{3}$  | $\sqrt{75}$ or $5\sqrt{3}$   | A1cso      |
|  |   |  |            |
| <b>(b)</b>   | $(x - 11)^2 + (y - 5\sqrt{3})^2 = 16$   | M1: $(x \pm 11)^2 + (y \pm \text{their } k)^2 = 4^2$<br><b>Equation must be of this form and must use <math>x</math> and <math>y</math> not other letters. <math>k</math> could be their last answer to part (a). Allow their <math>k \neq 0</math> or just the letter <math>k</math>.</b> | M1A1       |
|  |   | A1: $(x - 11)^2 + (y - 5\sqrt{3})^2 = 16$<br>or $(x - 11)^2 + (y - 5\sqrt{3})^2 = 4^2$<br><b>NB <math>5\sqrt{3}</math> must come from correct work in (a) and allow awrt 8.66</b>  |            |
|  | Allow in expanded form for the final A1<br>e.g. $x^2 - 22x + 121 + y^2 - 10\sqrt{3}y + 75 = 16$ |  | <b>[2]</b> |
| <b>Total 5</b>   |   |  |            |
| <b>Watch out for:</b>  |   |  |            |
| <div style="text-align: center;"> <math>(a) OQ = \sqrt{(6\sqrt{5})^2 + 4^2} = \sqrt{46}</math> M1<br/> <math>y_Q = \sqrt{46 - 11^2}</math> M0 (<math>OQ &lt; 11</math>)<br/> <math>y_Q = \sqrt{75}</math> A0<br/> <math>(b) (x - 11)^2 + (y - 5\sqrt{3})^2 = 16</math> M1A0                     </div> |   |  |            |



| Question Number       | Scheme  |  | Marks          |
|-----------------------|---|--|----------------|
| <p><b>10. (a)</b></p> | $\frac{1}{2}(9x + 6x)4x$ <p>or <math>2x \times 15x</math></p> <p>or <math>\left(\frac{1}{2}4x \times (9x - 6x) + 6x \times 4x\right)</math></p> <p>or <math>6x^2 + 24x^2</math></p> <p>or <math>\left(9x \times 4x - \frac{1}{2}4x \times (9x - 6x)\right)</math></p> <p>or <math>36x^2 - 6x^2</math></p>   | <p>M1: Correct attempt at the area of a trapezium.</p> <p>Note that <math>30x^2</math> on its own or <math>30x^2</math> from incorrect work e.g. <math>5x \times 6x</math> is M0.</p> <p>If there is a clear intention to find the area of the trapezium correctly allow the M1 but the A1 can be withheld if there are any slips.</p> | <p>M1A1cso</p> |
|                       | $\Rightarrow 30x^2 y = 9600 \Rightarrow y = \frac{9600}{30x^2} \Rightarrow y = \frac{320}{x^2} *$   | <p>A1: Correct proof with at least one intermediate step and no errors seen.</p> <p><b>“y =” is required.</b></p>  |                |
|                       |   |  |                |
| <p><b>(b)</b></p>     | $(S =) \frac{1}{2}(9x + 6x)4x + \frac{1}{2}(9x + 6x)4x + 6xy + 9xy + 5xy + 4xy$   |  | <p>M1A1</p>    |
|                       | <p>M1: An attempt to find the area of <b>six</b> faces of the prism. The 2 trapezia may be combined as <math>(9x + 6x)4x</math> or <math>60x^2</math> and the 4 other faces may be combined as <math>24xy</math> but all six faces must be included. There must be attempt at the areas of two trapezia that are dimensionally correct.</p> <p>A1: Correct expression in any form.</p> <p>Allow just <math>(S =) 60x^2 + 24xy</math> for M1A1</p> |  |                |
|                       | $y = \frac{320}{x^2} \Rightarrow (S =) 30x^2 + 30x^2 + 24x\left(\frac{320}{x^2}\right)$   |  | <p>M1</p>      |
|                       | <p>Substitutes <math>y = \frac{320}{x^2}</math> into their expression for <math>S</math> (may be done earlier). <math>S</math> should have at least one <math>x^2</math> term and one <math>xy</math> term but there may be other terms which may be dimensionally incorrect.</p>   |  |                |
|                       | <p>So, <math>(S =) 60x^2 + \frac{7680}{x} *</math></p>  | <p>Correct solution only.</p> <p>“S = “ is <b>not</b> required here.</p>   | <p>A1* cso</p> |
|                       |   | [4]  |                |

|   |  |  |            |
|---|--|--|------------|
| <b>10(c)</b>  | $\frac{dS}{dx} = 120x - 7680x^{-2} \left\{ = 120x - \frac{7680}{x^2} \right\}$                                   | M1: Either $60x^2 \rightarrow 120x$ or $\frac{7680}{x} \rightarrow \frac{\pm \lambda}{x^2}$  | M1         |
|   |  | A1: Correct differentiation (need not be simplified).  | A1 aef     |
|   | $120x - \frac{7680}{x^2} = 0$<br>$\Rightarrow x^3 = \frac{7680}{120}; = 64 \Rightarrow x = 4$                    | M1: $S' = 0$ and “their $x^3 = \pm$ value”<br>or “their $x^{-3} = \pm$ value” Setting their $\frac{dS}{dx} = 0$<br>and “candidate’s ft <b>correct</b> power of $x = a$ value”. <b>The power of <math>x</math> must be consistent with their differentiation.</b> If inequalities are used this mark cannot be gained until candidate states value of $x$ or $S$ from their $x$ without inequalities. $S' = 0$ can be implied by<br>$120x = \frac{7680}{x^2}$ . Some may spot that $x = 4$ gives<br>$S' = 0$ and provided they clearly show $S'(4) = 0$ allow this mark as long as $S'$ is correct. (If $S'$ is incorrect this method is allowed if their derivative is clearly zero for their value of $x$ ) | M1A1cso    |
|   |  | A1: $x = 4$ <b>only</b> ( $x^3 = 64 \Rightarrow x = \pm 4$ scores A0)<br>Note that the value of $x$ is not explicitly required so the use of $x = \sqrt[3]{64}$ to give $S = 2880$ would imply this mark.  |            |
|   | <b>Note some candidates stop here and do not go on to find <math>S</math> – maximum mark is 4/6</b>              |  |            |
| $\{x = 4,\}$<br>$S = 60(4)^2 + \frac{7680}{4} = 2880 \text{ (cm}^2\text{)}$ | Substitute candidate’s value of $x (\neq 0)$ into a formula for $S$ . <b>Dependent on both previous M marks.</b> | ddM1   |            |
|   | 2880 cso ( <b>Must come from correct work</b> )  | A1 cao and cso   |            |
|   |  |  | <b>[6]</b> |

|   |  |   |                 |
|---|--|---|-----------------|
| <b>10(d)</b>  | $\frac{d^2S}{dx^2} = 120 + \frac{15360}{x^3} > 0$ $\Rightarrow \text{Minimum}$   | <p>M1: Attempt <math>S'' (x^n \rightarrow x^{n-1})</math> <b>and</b> considers sign.<br/>                 This mark requires an attempt at the second derivative and <b>some consideration of its sign</b>.<br/>                 There does not necessarily need to be any substitution. An attempt to solve <math>S'' = 0</math> is M0</p>   | M1A1ft          |
|   |  | <p>A1: <math>120 + \frac{15360}{x^3}</math> and <math>&gt; 0</math> <b>and</b> conclusion.<br/>                 Requires a <b>correct</b> second derivative of <math>120 + \frac{15360}{x^3}</math> (need not be simplified) <b>and</b> a valid reason (e.g. <math>&gt; 0</math>), <b>and</b> conclusion.<br/>                 Only follow through a correct second derivative i.e. <math>x</math> may be incorrect <b>but must be positive</b> and/or <math>S''</math> may have been <u>evaluated</u> incorrectly.</p> |                 |
|   | <p><b>A correct <math>S''</math> followed by <math>S''("4") = "360"</math> therefore minimum would score no marks in (d)</b><br/> <b>A correct <math>S''</math> followed by <math>S''("4") = "360"</math> which is positive therefore minimum would score both marks</b></p> |   |                 |
|   |  |   | <b>[2]</b>      |
| <b>Note parts (c) and (d) can be marked together.</b> |  |   |                 |
|   |  |   | <b>Total 14</b> |

