





Question Number	Scheme	Marks
5c	First M1 for a complete method to find the speed / velocity (Could involve two <i>suvat</i> equations) Condone sign errors but must have correct numbers in their equation(s) First A1 for a correct equation (or equations) Second A1 for 16 or 16.3 (m s ⁻¹) Must be <i>positive (speed)</i>	
6a		B1 shape B1 270, V (2)
6b	$\frac{V}{0.6} = \frac{5V}{3}$ Given answer	M1A1 (2)
6c	Time decelerating is $5V$	B1
	$\frac{1}{2}V\frac{5V}{3} + (270 - 5V - \frac{5V}{3})V + \frac{1}{2}V \cdot 5V = 1500$	M1 A2
	OR: $\frac{1}{2}(270 + 270 - 5V - \frac{5V}{3})V = 1500$	
	$V^2 - 81V + 450 = 0$ Given answer	DM1A1 (6)
6d	$V^2 - 81V + 450 = 0$ or $V = \frac{81 \pm \sqrt{81^2 - 4 \times 450}}{2}$	M1 solving
	$(V - 6)(V - 75) = 0$	A1 A1
	$V = 6$ since $(5 \times 75) > 270$ or $V = 75$ unrealistic	B1 (4)
		14
	Notes	
6a	First B1 for a trapezium with line starting at the origin Second B1 for 270 and V correctly marked	
6b	M1 for $(t =) \frac{V}{0.6}$; N.B. M1A0 for $V=0.6t$ then answer Must see division or intermediate step from $V=0.6t$ e.g. Changing 0.6 into 3/5. A1 for $t = \frac{5V}{3}$ Given answer	


Question Number	Scheme	Marks
6c	<p>B1 for $5V$ identified appropriately</p> <p>First M1 for clear attempt to equate the <i>total</i> area under graph to 1500.</p> <p>(Must include all 3 parts (if not using the trapezium rule) with $\frac{1}{2}$ seen at least once to give equation in V only; may use (1 triangle + 1 trapezium) or (rectangle - trapezium)</p> <p>(May use <i>suvat</i> for one or more parts of the area)</p> <p>A2 for a correct equation, -1 e.e.o.o.</p> <p>Second DM1 dependent on first M1 for multiplying out and collecting terms and putting into appropriate form</p> <p>Third A1 for correct equation. Given answer</p>	
6d	<p>First M1 for solving their 3 term quadratic equation for V</p> <p>N.B. This M1 can be implied by two correct roots but if either answer incorrect then an explicit method must be shown for this M mark.</p> <p>First A1 for $V = 6$</p> <p>Second A1 for $V = 75$</p> <p>B1 on ePEN but treat as DM1, dependent on both previous A marks, for either reason</p>	
7a	$T - 3mg\sin\alpha - F = 3ma$	M1A1
	$4mg - T = 4ma$	M1A1 (4)
7b	$F = \frac{1}{4}R; R = 3mg\cos\alpha$	B1; M1A1
	$T - 2.4mg = 3ma$ $4mg - T = 4ma$	M1
	$a = \frac{8g}{35} \quad \text{Given answer}$	A1 (5)
7c	Particles have same acceleration	B1 (1)
7d	$v^2 = 2 \times \frac{8g}{35} \times 1.75 \quad (= 0.8g)$	M1 A1
	$-3mg\sin\alpha - F = 3ma'$	M1
	$a' = -0.8g$	A1
	$0 = 0.8g + 2 \times (-0.8g)s$	M1 A1
	Total distance = $0.5 + 1.75 = 2.25$ (m) Accept 2.3 (m)	A1 (7)
		17
	Notes	
7a	<p>First M1 for equation of motion for A with usual rules</p> <p>First A1 for a correct equation</p> <p>Second M1 for equation of motion for B with usual rules</p> <p>Second A1 for a correct equation</p> <p>N.B. If using different tension in second equation, M0 for that equation</p>	


Question Number	Scheme	Marks
7.(a)		<p>B1 shape</p> <p>B1 figs. (V,T,180) (2)</p>
(b)	<p>Time accelerating = $V/1 = V$</p> <p>Time decelerating = $V/0.5 = 2V$</p> <p>Time at constant speed, $T = 180 - (2V + V)$ $T = 180 - 3V$ Printed answer</p>	<p>M1</p> <p>A1 (2)</p>
(c)	$\frac{1}{2}(180 + 180 - 3V)V = 4800$ $V^2 - 120V + 3200 = 0$ $(V - 40)(V - 80) = 0$ $V = 40 \text{ or } 80 \text{ or both, since } (180 - 3 \times 80) < 0$	<p>M1 A1 A1</p> <p>A1</p> <p>DM1 A1, M1 (7) 11</p>
Notes		
7.(a)	<p>First B1 for a trapezium, starting at the origin and finishing on the t-axis. Second B1 for V, T with delineators or marked on the top of the trapezium or oe and 180 correctly positioned.</p>	
(b)	<p>M1 for both Time accelerating = $V/1 = V$ and Time decelerating = $V/0.5 = 2V$ M0 if no working for the $2V$ as it's a 'Show that' or if they use $V/-0.5$ and fudge the -ve sign A1 for $T = 180 - (2V + V) = 180 - 3V$ Printed answer</p>	

Question	Scheme	Marks	Notes
5. (a)		B1 B1 B1	One graph correct shape Both graphs correct shape, on same sketch and intersecting (with different start times) Figs 10,20,25,40 shown (with 20 as the second start time) Ignore all vertical lines
		(3)	
(b)	$20 + 10$	M1	Complete method
	$= 30$	A1	
		(2)	
(c)	$\frac{40}{t_1 - 20} = \frac{25}{10}$	M1	Complete method to find time when Q reaches 40 m s^{-1}
		A1	Correct unsimplified equation
	$\Rightarrow t_1 = 36$	A1	
Or:	Time to reach 40 m s^{-1} is $\frac{40}{2.5} (= 16)$ (M1A1)		
	Time from start $= \frac{40}{2.5} + 20 = 36$ (A1)		(seen or implied)
		M1	Find distance travelled by either train at $t = T$
	$\frac{(T + T - 10)}{2} \times 25$	A1	One correct
	$\frac{(T - 20 + T - 36)}{2} \times 40$	A1ft	Both correct. Follow their 36
	Equate and solve for T	dM1	
	$T = 66\frac{1}{3}$	A1	Accept 66 or better
		(8)	
		13	


Question Number	Scheme	Marks	Notes
6.(a)	$v = u + at \Rightarrow 14 = 3.5a$	M1	Use of <i>suvat</i> to form an equation in a
	$a = 4$	A1(2)	
6.(b)		B1	Graph for A or B
		B1	Second graph correct and both graphs extending beyond the point of intersection
		B1	Values 3.5, 14, T shown on axes, with T not at the point of intersection. Accept labels with delineators.
		(3)	NB 2 separate diagrams scores max B1B0B1
6.(c)	$\frac{1}{2}T \cdot 3T$, $\frac{(T+T-3.5)}{2} \cdot 14$	M1	Find distance for A or B in terms of T only. Correct area formulae: must see $\frac{1}{2}$ in area formula and be adding in trapezium
		A1	One distance correct
		A1	Both distances correct
	$\frac{1}{2}T \cdot 3T = \frac{(T+T-3.5)}{2} \cdot 14$ $\frac{1}{2}T \cdot 3T = \frac{1}{2} \times 4 \times 3.5^2 + 14(T-3.5)$	M1	Equate distances and simplify to a 3 term quadratic in T in the form $aT^2 + bT + c = 0$
	$3T^2 - 28T + 49 = 0$	A1	Correct quadratic
	$(3T-7)(T-7) = 0$	M1	Solve 3 term quadratic for T
	$T = \frac{7}{3}$ or 7	A1	Correct solution(s) - can be implied if only ever see $T = 7$ from correct work.
	but $T > 3.5$, $T = 7$	A1 (8)	
6.(d)	73.5 m	B1 (1)	From correct work only. B0 if extra answers.
6.(e)		B1	(A) Condone missing 4
		B1	(B) Condone graph going beyond $T = 7$ Must go beyond 3.5. Condone no 3.
		B1 (3)	(A) Condone graph going beyond $T = 7$ Must go beyond 3.5. B0 if see a <u>solid</u> vertical line. Sometimes very difficult to see. If you think it is there, give the mark.
		[17]	Condone separate diagrams. See next page

	<p>Alternative for (c) for candidates with a sketch like this:</p> 	<p>B1 B1 B0</p>	<p>Treat as a special case. B1B1B0 on the graph and then max 5/8 for (c) if they do not solve for the T in the question.</p>
	$\frac{1}{2} \times 3 \times (T + 3.5)^2 = \frac{1}{2} \times 4 \times 3.5^2 + 14T$	<p>M1</p>	<p>Use diagram to find area</p>
		<p>A1</p>	<p>One distance correct</p>
		<p>A1</p>	<p>Both distances correct</p>
	$12T^2 - 28T - 49 = 0$	<p>M1</p>	<p>Simplify to a 3 term quadratic in T</p>
		<p>A1</p>	<p>Correct quadratic</p>
	$(2T - 7)(6T + 7) = 0$	<p>M1</p>	<p>Complete method to solve for the T in the question</p>
	$T = \frac{7}{2} \text{ or } -\frac{7}{6}$	<p>A1</p>	<p>Correct solution(s) - can be implied if only ever see Total = 7</p>
	<p>Total time = 7</p>	<p>A1 (8)</p>	

Question Number	Scheme	Marks	Notes
6(a)	$v_1 = 8 \times 1.5 (= 12)$ $v_2 = 12 + 0.8 \times 20$ $v_2 = 28 \text{ m s}^{-1}$	M1 M1 A1 (3)	Use of $v = u + at$ or equivalent for $t = 8$ Follow their 12
(b)		B1 B1ft	shape nos: 8,28; 12,28 indicated. Follow their 12, 28
(c)	first 8 s: $\text{dist} = \frac{1}{2} \times 8 \times 12 (= 48)$ next 20 s: $\text{dist} = \frac{1}{2} \times (12 + 28) \times 20 (= 400)$ Total dist = 448 m	(2) M1 A1ft A1ft A1 (4)	Correct method for distance for the triangle (0-8) or the trapezium (8-28) Follow their 12 Follow their 12, 28 Correct answer only (cao)
(d)	$0 = 28^2 - 2 \times 2.8s$ $s = \frac{28^2}{2 \times 2.8} (= 140)$ $448 + 140 + 28T = 2000$ $T = \frac{2000 - 448 - 140}{28} = 50.4$	M1 A1ft DM1 A1 (4) [13]	Find area of right hand triangle or an expression in T for the trapezium (rectangle + triangle). Follow their 28 Form an equation in T for their 16, 448 and 140 Or better (50.42857...) Accept 50.

Question Number	Scheme	Marks
5a		
	Basic shape 20, 4T and T placed correctly	B1 DB1
		(2)
5b	Use of $v = u + at$: constant speed = $0.6 \times 20 = 12 \text{ (ms}^{-1}\text{)}$ (Speed at end = $12 - 0.3T$)	M1A1
	Using v-t graph: Distance: $705 = \frac{12}{2}(4T + (20 + 4T)) + \frac{T}{2}(12 + (12 - 0.3T))$ $= 48T + 120 + 12T - 0.15T^2 = 60T + 120 - 0.15T^2$	M1A2
	Form 3 term quadratic and solve for T: $\Rightarrow 3T^2 - 1200T + 11700 = 0 \quad (T^2 - 400T + 3900 = 0)$	M1
	$\Rightarrow (T - 10)(T - 390) = 0 \quad T = 10 \text{ only}$	A1
		(7)
	Alternative:	
	Use of $v = u + at$: constant speed = $0.6 \times 20 = 12 \text{ (ms}^{-1}\text{)}$	M1A1
	Using $s = ut + \frac{1}{2}at^2$: $705 = (0.3 \times 400) + (4T \times 12) + (12T - 0.15T^2)$ M1A2	M1A2
	$\Rightarrow 0.15T^2 - 60T + 585 = 0 \quad (T^2 - 400T + 3900 = 0)$	
	$\Rightarrow (T - 10)(T - 390) = 0 \quad T = 10 \text{ only}$	M1A1
	(7)	
5c	Extra time: $(2 \times 20) - \text{their } T$ OR $\frac{12 - 0.3 \times \text{their } T}{0.3}$	B1
	Total time: $20 + 5T + 40 - T$ (their T)	M1
	$= 100 \text{ (s)}$	A1
		(3)
	Alternative: Total time to decelerate to rest = $12/0.3 = 40$	B1
	Total time A to C = $20 + 4T + 40 = 100$	M1A1
		[12]

Question Number	Scheme	Marks
	Notes for question 5	
5a	First B1 for basic shape. Allow if 'extra triangle' on end included, <u>provided B clearly marked</u>	
	Second DB1 : may use, 20, 20 + 4T, 20 + 5T	
5b	First M1 for attempt to find constant speed ($v = u + at$ or $a = \text{gradient}$) 20 x 0.6	
	First A1 for 12	
	Second (generous) M1 for clear attempt to use $705 = \text{total area under the graph}$ to give an equation in T only but must see $\frac{1}{2}$ used somewhere N.B. M0 if just a trapezium oe is used	
	Second A1 and Third A1: for any correct equation, -1 e.e.o.o.	
	Third M1 for forming and attempt to solve a 3 term quadratic (need <i>evidence</i> of solving e.g. formula or factorising, if T values are incorrect) otherwise this M mark can be implied if they state that $T = 10$ with no working. ($T = 390$ NOT needed)	
	Fourth A1 for $T = 10$.	
	N.B. For total area, could see: Trapezium + Rectangle + Triangle $705 = \frac{12}{2}(4T + (20 + 4T)) + T(12 - 0.3T) + \frac{1}{2}T \times 0.3T$ Triangle + Rectangle + Trapezium $705 = \frac{1}{2}.20.12 + (4T \times 12) + \frac{1}{2}T(12 + 12 - 0.3T)$ Triangle + Rectangle + Rectangle + Triangle $705 = \frac{1}{2}.20.12 + (4T \times 12) + T(12 - 0.3T) + \frac{1}{2}T \times 0.3T$ Triangle + Rectangle + Trapezium (at top) $705 = \frac{1}{2}.20.12 + 5T(12 - 0.3T) + \frac{1}{2}0.3T(5T + 4T)$ Rectangle - triangle- triangle $705 = 12(20 + 5T) - \frac{1}{2}.20.12 - \frac{1}{2}T \times 0.3T$	
5c	B1 for either additional time is $\frac{12}{0.3} - T$ or time to decelerate is $\frac{12}{0.3}$	
	M1 for a correct method to find the total time, using <i>their T</i> $= 20 + 4T + T + \frac{12}{0.3} - T$ or $20 + 4T + \frac{12}{0.3}$	
	A1 for 100 cao	

Question Number	Scheme	Marks
<p>4(a)</p> <p>ALT</p>	$0^2 = 11.2^2 - 2gd$ $d = 6.4$ $\text{max ht.} = 3.6 + 6.4 = 10 \text{ m}$ $11.2^2 = u^2 - 2g \times 3.6$ $u = 14$ $0^2 = 14^2 - 2gh$ $h = 10 \text{ m}$	<p>M1 A1</p> <p>A1</p> <p>A1</p> <p>(4)</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>(4)</p>
(b)	$10 = \frac{1}{2}gt^2$ $t = \frac{10}{7}$ $\text{Total} = 2 \times \frac{10}{7} = 2.9 \text{ or } 2.86$	<p>M1 A1</p> <p>A1</p> <p>dM1 A1</p> <p>(5)</p>
(c)		<p>B1 single line</p> <p>dB1 $V < -11.2$</p> <p>B1 11.2</p> <p>B1 1.1(4)</p> <p>(4)</p> <p>13</p>
Notes		
<p>4(a)</p> <p>ALT</p>	<p>M1 for a complete method to find d ($d =$ distance from A to top) First A1 for a correct equation in d only. Second A1 for $d = 6.4$ Third A1 for $6.4 + 3.6 = 10$ (m)</p> <p>M1 for a complete method (must have 2nd equation) to find h First A1 for $u = 14$ Second A1 for correct 2nd equation Third A1 for $h = 10$ (m)</p>	
4(b)	<p>First M1 for a complete method to find an intermediate time (A to top or A to O) First A1 for a correct equation or equations. Second A1 for any intermediate time (e.g. $At_{\text{TOP}} = 8/7$, $AtO = 2/7$, $AtO = 18/7$, $AtA =$</p>	

Question Number	Scheme	Marks
8. (a)	<p>The graph shows velocity (v) on the vertical axis and time (t) on the horizontal axis. The horizontal axis has markings at 0, 30, 90, 150, and 180. A trapezium is formed by the points (0,0), (30,20), (150,20), and (180,0). A triangle is formed by the points (0,0), (90,40), and (180,0).</p>	B1 trapezium B1 triangle & overlap B1 figs (3)
(b)	$\frac{1}{2}(90+60).20 = 1500$ $1500 = \frac{1}{2}a.90^2$ $a = \frac{10}{27} \text{ ms}^{-1} \text{ or decimal}$	M1 A1 M1 A1 ft A1 (5)
(c)	$\frac{10t}{27} = 20$ $t = 54 \text{ s}$ $t = 126 \text{ s}$	M1 A1 A1 A1 ft (4)
(d)	$\frac{10}{27} \times 90 \left(= \frac{100}{3} \right)$ $\frac{100}{3} \times 6 - \frac{1}{2} \cdot \frac{10}{27} \cdot 6^2 \left(= \frac{580}{3} \right)$ $d = \frac{580}{3} - (20 \times 6)$ $= \frac{220}{3} \text{ m or decimal}$	M1 DM1 A1 DM1 A1 (5) 17