



1. [In this question  $\mathbf{i}$  and  $\mathbf{j}$  are perpendicular unit vectors in a horizontal plane.]

A particle  $P$  moves in such a way that its velocity  $\mathbf{v}$  m s<sup>-1</sup> at time  $t$  seconds is given by

$$\mathbf{v} = (3t^2 - 1)\mathbf{i} + (4t - t^2)\mathbf{j}$$

(a) Find the magnitude of the acceleration of  $P$  when  $t = 1$

**(5)**

Given that, when  $t = 0$ , the position vector of  $P$  is  $\mathbf{i}$  metres,

(b) find the position vector of  $P$  when  $t = 3$

**(5)**

Lined area for student answers.



Leave blank

**Question 1 continued**

Handwriting practice area consisting of 31 horizontal lines.

**(Total 10 marks)**

Q1

Two empty boxes for marking.



P 4 0 6 9 0 A 0 3 2 4





3.




Figure 1

A uniform rod  $AB$ , of mass 5 kg and length 4 m, has its end  $A$  smoothly hinged at a fixed point. The rod is held in equilibrium at an angle of  $25^\circ$  above the horizontal by a force of magnitude  $F$  newtons applied to its end  $B$ . The force acts in the vertical plane containing the rod and in a direction which makes an angle of  $40^\circ$  with the rod, as shown in Figure 1.

(a) Find the value of  $F$ . (4)

(b) Find the magnitude and direction of the vertical component of the force acting on the rod at  $A$ . (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Leave blank

**Question 3 continued**

[This section contains 28 horizontal lines for writing the answer to Question 3 continued.]


**(Total 8 marks)**

**Q3**



P 4 0 6 9 0 A 0 7 2 4

4.



**Figure 2**

A uniform circular disc has centre  $O$  and radius  $4a$ . The lines  $PQ$  and  $ST$  are perpendicular diameters of the disc. A circular hole of radius  $2a$  is made in the disc, with the centre of the hole at the point  $R$  on  $OP$  where  $OR = 2a$ , to form the lamina  $L$ , shown shaded in Figure 2.

(a) Show that the distance of the centre of mass of  $L$  from  $P$  is  $\frac{14a}{3}$ . (4)

The mass of  $L$  is  $m$  and a particle of mass  $km$  is now fixed to  $L$  at the point  $P$ . The system is now suspended from the point  $S$  and hangs freely in equilibrium. The diameter  $ST$  makes an angle  $\alpha$  with the downward vertical through  $S$ , where  $\tan \alpha = \frac{5}{6}$ .

(b) Find the value of  $k$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---











5.




Figure 3

A small ball  $B$  of mass  $0.25$  kg is moving in a straight line with speed  $30 \text{ m s}^{-1}$  on a smooth horizontal plane when it is given an impulse. The impulse has magnitude  $12.5 \text{ N s}$  and is applied in a horizontal direction making an angle of  $(90^\circ + \alpha)$ , where  $\tan \alpha = \frac{3}{4}$ , with the initial direction of motion of the ball, as shown in Figure 3.

- (i) Find the speed of  $B$  immediately after the impulse is applied.
- (ii) Find the direction of motion of  $B$  immediately after the impulse is applied.

(6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









- 6. A car of mass 1200 kg pulls a trailer of mass 400 kg up a straight road which is inclined to the horizontal at an angle  $\alpha$ , where  $\sin \alpha = \frac{1}{14}$ . The trailer is attached to the car by a light inextensible towbar which is parallel to the road. The car's engine works at a constant rate of 60 kW. The non-gravitational resistances to motion are constant and of magnitude 1000 N on the car and 200 N on the trailer.

At a given instant, the car is moving at  $10 \text{ m s}^{-1}$ . Find

- (a) the acceleration of the car at this instant, (5)

- (b) the tension in the towbar at this instant. (4)

The towbar breaks when the car is moving at  $12 \text{ m s}^{-1}$ .

- (c) Find, using the work-energy principle, the further distance that the trailer travels before coming instantaneously to rest. (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---











7.




Figure 4

A small stone is projected from a point  $O$  at the top of a vertical cliff  $OA$ . The point  $O$  is  $52.5$  m above the sea. The stone rises to a maximum height of  $10$  m above the level of  $O$  before hitting the sea at the point  $B$ , where  $AB = 50$  m, as shown in Figure 4. The stone is modelled as a particle moving freely under gravity.

- (a) Show that the vertical component of the velocity of projection of the stone is  $14 \text{ m s}^{-1}$ . (3)
  
- (b) Find the speed of projection. (9)
  
- (c) Find the time after projection when the stone is moving parallel to  $OB$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---







**Question 7 continued**

A series of horizontal lines for writing, intended for the continuation of Question 7.



Leave  
blank

**Question 7 continued**

Lined area for writing answers.

**(Total 17 marks)**

**Q7**

--	--

**TOTAL FOR PAPER: 75 MARKS**

**END**

